

The FP_m -Conjecture for a Class of Metabelian Groups

Dessislava H. Kochloukova

DPMMS, University of Cambridge, 16 Mill Lane, Cambridge CB2 1SB, England

Communicated by Jan Saxl

Metadata, citation and similar papers at core.ac.uk

We prove that if $A \rightarrow G \rightarrow Q$ is a short exact sequence of groups where G is finitely generated, A and Q are abelian, A is a \mathbb{Z} -torsion Krull dimension one $\mathbb{Z}Q$ -module via conjugation then the FP_m -Conjecture holds. In general if G is of type FP_m and either the extension is split or A is \mathbb{Z} -torsion we show that A is m -tame as $\mathbb{Z}Q$ -module. © 1996 Academic Press, Inc.

1. INTRODUCTION

Let G be a finitely generated metabelian group with an abelian normal subgroup A and an abelian quotient $Q \simeq G/A$. Then A is a finitely generated (left) $\mathbb{Z}Q$ -module, where Q acts via conjugation. One geometrical invariant of the $\mathbb{Z}Q$ -module A is

$$\Sigma_A(Q) = \{ \chi: Q \rightarrow \mathbb{R} \mid \chi \neq 0, A \text{ is finitely generated as } \mathbb{Z}Q_\chi\text{-module} \} / \sim,$$

where $Q_\chi = \{ q \in Q: \chi(q) \geq 0 \}$ and \sim is an equivalence relation in $\text{Hom}_{\mathbb{Z}}(Q, \mathbb{R})$ with equivalence classes $\mathbb{R}^+ \chi$, where $\chi \in \text{Hom}_{\mathbb{Z}}(Q, \mathbb{R})$. If n is the rank of Q then $\Sigma_A(Q)$ can be viewed as a subset of the unit sphere S^{n-1} in $\mathbb{R}^n \simeq \text{Hom}_{\mathbb{Z}}(Q, \mathbb{R})$. This presentation of $\Sigma_A(Q)$ as a subset of S^{n-1} depends on the chosen isomorphism between \mathbb{R}^n and $\text{Hom}_{\mathbb{Z}}(Q, \mathbb{R})$. We assume from now on that such an isomorphism is fixed and consider $\Sigma_A(Q)$ as a subset of S^{n-1} .

The $\mathbb{Z}Q$ -module A is said to be m -tame if any characters χ_1, \dots, χ_m lying in

$$\begin{aligned} \Delta_A(Q) &= \{ \chi: Q \rightarrow \mathbb{R} \mid \chi \neq 0, A \text{ is not finitely generated as } \mathbb{Z}Q_\chi\text{-module} \} \\ &\subseteq \mathbb{R}^n \end{aligned}$$

are contained in an open half subspace of $\text{Hom}_{\mathbb{Z}}(Q, \mathbb{R}) \simeq \mathbb{R}^n$. The class of 2-tame $\mathbb{Z}Q$ -modules was first considered by Bieri and Strebel in [5].

A group G is said to be of type FP_m if the trivial $\mathbb{Z}G$ -module \mathbb{Z} has a projective resolution

$$\cdots \rightarrow P_i \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \mathbb{Z}$$

with P_i finitely generated as $\mathbb{Z}G$ -module for $i \leq m$. In [5] Bieri and Strebel prove that A is 2-tame if and only if G is finitely presented and show that this happens precisely when G is of type FP_2 . Bieri and Groves [3] show that if G is of type FP_m and A is of exponent p for some prime integer p then A is m -tame as $\mathbb{Z}Q$ -module. In [2] Bieri suggests the following conjecture

FP_m -Conjecture. G is of type FP_m if and only if A is m -tame.

In the case where G is a finitely generated metabelian group of finite Prufer rank, i.e., there is an integer d such that every finitely generated subgroup may be generated by at most d -generators, the FP_m -Conjecture is proved by Åberg [1]. In [11] Noskov generalizes Åberg's proof and shows that if A is torsion-free, G is a split extension of A by Q , and G is of type FP_m then A is m -tame.

One of the main results obtained in the paper is the following theorem.

THEOREM A. *Let G be a finitely generated group with a normal subgroup A and an abelian quotient $Q \simeq G/A$ such that A is a torsion abelian group and has Krull dimension one as $\mathbb{Z}Q$ -module. If A is m -tame as $\mathbb{Z}Q$ -module then G is of type FP_m .*

In order to prove Theorem A we construct a CW -complex Y , acted on by G with the following properties:

- (i) Y/G is a compact space;
- (ii) Y is $(m - 1)$ -acyclic;
- (iii) the stabilizers in G of cells in Y are polycyclic.

Then we can apply the following criterion due to K. Brown.

THEOREM [9]. *Suppose G is a group acting on $(m - 1)$ -acyclic CW -complex Y . Let D be a directed index set and let $\{Y_\alpha\}_{\alpha \in D}$ be a filtration of Y by G -subcomplexes Y_α such that Y_α has finite m -skeleton mod G for all $\alpha \in D$. Furthermore, assume that every stabilizer G_e , where e is an i -cell of Y , is of type FP_{m-i} , $i \leq m$. Then G is of type FP_m if and only if the filtration $\{Y_\alpha\}_{\alpha \in D}$ is essentially $(m - 1)$ -acyclic (i.e., for every $\alpha \in D$ there exists $\beta \in D$, $\alpha \leq \beta$ such that $H_i(Y_\alpha) \rightarrow H_i(Y_\beta)$ is the trivial homomorphism for each $i \leq m - 1$).*

Remark. In the particular case when the set D contains only one element $Y = Y_\alpha$ and the stabilizers are polycyclic, this being exactly the situation we will meet later, Åberg notices in [1] that if Y is $(m - 1)$ -acyclic then G is of type FP_m .

The construction of the CW -complex Y resembles the construction given in [1] but we do not require that G be a split extension of A by Q . The complex Y is defined as a subspace of the product X of some G -trees X_v and the G -trees X_v are similar to the valuation trees considered in the split case in [1]. The tree X_v is the tree associated to a Cayley graph of G and a discrete character v of Q . Details about this construction can be found in [7]; in 2.1 we explain all the properties of X_v we will need later.

We generalize one of the main ideas of the proof of the FP_m -Conjecture for metabelian groups of finite Prüfer rank given by Åberg [1] and prove the following theorem

THEOREM B. *Let $A \rightarrow G \rightarrow Q$ be a short exact sequence of groups with G finitely generated, A and Q abelian, and either the extension is split or A is a torsion group. If G is of type FP_m then A is m -tame as $\mathbb{Z}Q$ -module.*

As a consequence of Theorems A and B we obtain

COROLLARY C. *If G is a finitely generated group which is an extension of a \mathbb{Z} -torsion Krull dimension one $\mathbb{Z}Q$ -module A by an abelian group Q then the FP_m -Conjecture holds.*

Recently K. U. Bux [10] has proved Corollary C in the case where A is an S -arithmetic ring of prime characteristic and the extension is split. He uses methods from algebraic number theory to prove that the group G acts cocompactly on a space of Åberg's type.

The paper is structured as follows: in 2.1 we define the G -tree X_v for a discrete character v of Q , in 2.2 we consider a complex of Åberg's type Y , the set of characters needed in the definition of the space Y is described in 2.3, in 2.4 we show that Y/G is a compact space, and finally in 2.5 we prove that the stabilizers in G of cells in Y are polycyclic.

The proof of Theorem B is discussed in Section 3. In 3.2 and 3.3 we define an element of $H^0(Q', (\mathbb{Z}G^{\mathbb{N}})_A)$ and in the case where A is not m -tame we show in 3.4 that the image of this element in $H_0(Q'', (\mathbb{Z}G^{\mathbb{N}})_A)$ is non-trivial, where Q' and Q'' are subgroups of Q with $Q' \times Q'' \simeq Q$. The last will contradict the fact that G is of type FP_m .

Since the FP_m -Conjecture holds for a group G if and only if it holds for a subgroup of finite index in G , we assume from now on that $Q \simeq G/A$ is free abelian of rank n . Therefore $Q = \mathbb{Z}^n \subset \mathbb{R}^n$, and from now on we consider the vector space \mathbb{R}^n equipped with the standard inner product.

2. PROOF OF THEOREM A

2.1. Trees Deriving from Cayley Graphs

Let G be a finitely generated metabelian group with an abelian normal subgroup A and an abelian quotient $Q \simeq G/A \simeq \mathbb{Z}^n$. Let v be a discrete character of Q with $v(Q) = \mathbb{Z}$. It is shown in [7] that to every Cayley graph Γ corresponding to a finite set of generators of G there can be assigned a tree Γ_v , which depends on the character v . Here we give a detailed construction of a tree X_v that is in fact Γ_v for a suitably chosen set of generators of G and the corresponding Cayley graph Γ .

Set $Q_v = \{q \in Q : v(q) \geq 0\}$ and let $q_{v,1}, \dots, q_{v,n-1}$ be generators of $\text{Ker } v$. Assume q_v is an element of Q such that $v(q_v) = 1$. Let A be generated as $\mathbb{Z}Q$ -module by the elements a_1, \dots, a_s and let π be a lifting of the projection map $G \rightarrow G/A$. We define A_v to be the $\mathbb{Z}Q_v$ -submodule of A generated by a_1, \dots, a_s and G_v to be the subgroup of G generated by $q_v^{\beta_v} A_v$ and $\pi(q_{v,i})$ for $1 \leq i \leq n-1$, where β_v is a negative integer such that all the commutators of $\pi(q_v)^{\pm 1}$, $\pi(q_{v,i})^{\pm 1}$ for $1 \leq i \leq n-1$ lie in $q_v^{\beta_v} A_v$. Note that if A is finitely generated as $\mathbb{Z}Q_v$ -module then by [5, Proposition 2.1] $A_v = A$, and if A is not finitely generated as $\mathbb{Z}Q_v$ -module then $A_v \neq A$ and $\bigcup_{z \in \mathbb{Z}} q_v^z A_v = A$.

Since $[\pi(q_{v,j})^{\pm 1}, \pi(q_{v,i})^{\pm 1}] \in q_v^{\beta_v} A_v$ we have $G_v \cap A = q_v^{\beta_v} A_v$. At the same time $\pi(q_v)\pi(q_{v,i})^{\pm 1}\pi(q_v)^{-1} \in q_v^{\beta_v} A_v \pi(q_{v,i})^{\pm 1} \subseteq G_v$ and for $a \in q_v^{\beta_v} A_v$ we have $\pi(q_v)a\pi(q_v)^{-1} = q_v a \in q_v^{\beta_v+1} A_v \subseteq q_v^{\beta_v} A_v$. So we get

$$\pi(q_v)G_v\pi(q_v)^{-1} \subseteq G_v. \quad (1)$$

We define G_v^+ to be the submonoid of G generated by G_v and $\pi(q_v)$. By (1)

$$G_v^+ = \bigcup_{z \geq 0} G_v \pi(q_v)^z. \quad (2)$$

Let T_0 be the G -set G/G_v with partial order given by $gG_v \leq hG_v$ if and only if $g^{-1}h \in G_v^+$. The character $\alpha_v: G \rightarrow \mathbb{Z}$ which extends the character v factors through a map $\chi_v: T_0 \rightarrow \mathbb{Z}$.

We call two points α, β of T_0 neighbours if $\alpha \leq \beta$, $\chi_v(\beta) - \chi_v(\alpha) = 1$, or $\beta \leq \alpha$, $\chi_v(\alpha) - \chi_v(\beta) = 1$. We link every two neighbours $\alpha \leq \beta$ with a closed unit interval and write $(1-t)\alpha + t\beta$ for the point of this interval having distances t and $1-t$ to α and β , respectively. The constructed intervals intersect only at their ends or do not intersect. We get a graph T with vertices T_0 and edges the constructed unit intervals between neighbours of T_0 . We extend $\chi_v: T_0 \rightarrow \mathbb{Z}$ to an \mathbb{R} -linear map $\chi_v: T \rightarrow \mathbb{R}$; in other words $\chi_v((1-t)\alpha + t\beta) = (1-t)\chi_v(\alpha) + t\chi_v(\beta) = \chi_v(\alpha) + t$ where $\alpha \leq \beta$ and $\chi_v(\beta) - \chi_v(\alpha) = 1$.

If $\alpha \leq \beta$ are neighbours then their images $g * \alpha$ and $g * \beta$ under the action of an element g of G are neighbours as well and $g * \alpha \leq g * \beta$. We make G act on T by $g * ((1 - t)\alpha + t\beta) = (1 - t)(g * \alpha) + t(g * \beta)$ using the G -action on T_0 . We write X_v for the graph T .

For any $g \in G$ we define $L_{g,v}$ to be the line in X_v spanned by the vertices $\{g\pi(q_v)^z G_v : z \in \mathbb{Z}\}$. Note that for any integer z there is an edge in X_v from $g\pi(q_v)^z G_v$ to $g\pi(q_v)^{z+1} G_v$. The restriction $\chi_v : L_{g,v} \rightarrow \mathbb{R}$ is a bijection. Since $G = \bigcup_{z \in \mathbb{Z}} A\pi(q_v)^z G_v$ and the edge between $g\pi(q_v)^z G_v$ and $g\pi(q_v)^{z+1} G_v$ is the unique one going to $g\pi(q_v)^{z+1} G_v$ (i.e., if $h_1 G_v < h_2 G_v$ are neighbours then $h_1 G_v = h_2 \pi(q_v)^{-1} G_v$) we get

$$X_v = \bigcup_{a \in A} L_{a,v}.$$

We write $[(a, r)]$ where $a \in A$ and $r \in \mathbb{R}$ for the unique element of $L_{a,v}$ whose image under χ_v is r . By the definition of the G -action on X_v we have $b * [(a, r)] = [(b + a, r)]$ where b, a are any elements of A and r is any real number.

Remark. The above description of X_v resembles the valuation tree associated to G and a valuation of A , extending the character v , in the case where A is an integral domain and 1-generated $\mathbb{Z}Q$ -module, and G is a split extension of A by Q [1, Chap. 3.2].

LEMMA 2.1. (i) For every $a, b \in A$ we have $L_{a,v} \cap L_{b,v} = \{[(a, r)] : r \leq \text{some integer depending on } a, b\}$ or $L_{a,v} = L_{b,v}$. In particular X_v is a tree.

(ii) The graph X_v can be represented as a union of ascending rays T_i , $i \geq 2$ (i.e., $\chi_v : T_i \rightarrow \mathbb{R}$ is injective and the image of this map is $[r_i, \infty)$ for some real number r_i) and a line T_1 (where $\chi_v : T_1 \rightarrow \mathbb{R}$ is a bijection) such that for every j we have $(\bigcup_{i \leq j} T_i) \cap T_{j+1} = \{w_j\}$ for a vertex w_j of X_v and $\chi_v(T_{j+1}) = [\chi_v(w_j), \infty)$.

Proof. (i) If a, b are elements of A and z is an integer then $[(a, z)] = [(b, z)]$ if and only if $q_v^{-z}(a - b) \in G_v \cap A = q_v^{\beta_v} A_v$. Let $z_0 = \sup\{z \in \mathbb{Z} : a - b \in q_v^{z+\beta_v} A_v\}$. If $z_0 = \infty$ we have $L_{a,v} = L_{b,v}$ and if $z \leq z_0 < \infty$ then $a - b \in q_v^{z_0+\beta_v} A_v \subseteq q_v^{z+\beta_v} A_v$. Thus $L_{a,v} \cap L_{b,v} = \{[(a, r)] : r \leq z_0\}$.

(ii) Since $\chi_v : L_{a,v} \rightarrow \mathbb{R}$ is a bijection for every element a of A , (ii) follows from (i).

2.2. A Complex of Aberg's Type

Let V be a finite set of characters of Q with value groups, the group of rational integers \mathbb{Z} , and for every character v in V let X_v be the tree defined in the previous section with the additional property that all β_v , needed in the definition of G_v , are equal to a negative integer β . Later we

will put some more restrictions on an upper bound of β in order to prove that the complex of Åberg's type constructed below has all the properties required in the Introduction.

We define a space $X = \prod_{v \in V} X_v$ and a map $f: X \rightarrow \prod_{v \in V} \mathbb{R}_v$ (where $\mathbb{R}_v = \mathbb{R}$ for every $v \in V$) given by $f(\prod_{v \in V} [(a_v, r_v)]) = \prod_{v \in V} r_v$ where a_v lies in A and r_v is a real number. Let $\chi: Q \otimes \mathbb{R} \rightarrow \prod_{v \in V} \mathbb{R}_v$ be the \mathbb{R} -linear map extending the map sending an element q of Q to $\prod_{v \in V} v(q)$. Denote Y the subspace $\{x \in X: f(x) \in \text{Im } \chi\}$ of X . If all X_v are valuation trees this construction is due to Åberg [1]. We make G act on X via the diagonal action (i.e., $g * \prod_{v \in V} [(a_v, r_v)] = \prod_{v \in V} g * [(a_v, r_v)]$).

LEMMA 2.2. *The space Y is invariant under this G -action.*

Proof. Let $\prod_{v \in V} [(a_v, r_v)]$ be a point of Y and let g be an element of G . Then $g * \prod_{v \in V} [(a_v, r_v)] = \prod_{v \in V} [(a'_v, r_v + \alpha_v(g))]$ where $ga_v \pi(q_v)^{[r_v]} G_v = a'_v \pi(q_v)^{[r_v] + \alpha_v(g)} G_v$ for some element a'_v of A and where $[r_v]$ is the rational integer such that $[r_v] - r_v \in [0, 1)$. We remind the reader that $\alpha_v: G \rightarrow \mathbb{Z}$ is the group homomorphism extending the character v . Since $\prod_{v \in V} \alpha_v(g) \in \text{Im } \chi$ we see that Y is invariant under G -action.

All the constructions considered up to now are quite general but from now on we assume that $M := \mathbb{Z}Q / \text{ann}_{\mathbb{Z}Q} A$ is a \mathbb{Z} -torsion ring of Krull dimension one. By [6] $\Sigma_M^c(Q) = \bigcup_P \Sigma_{M/P}^c(Q)$ where P runs through all minimal prime ideals of M and $\Sigma_-^c(Q)$ denotes the complement of $\Sigma_-(Q)$ in S^{n-1} . By [4] $\Sigma_{M/P}^c(Q)$ is the image of the projection to the unit sphere S^{n-1} in \mathbb{R}^n of a rationally defined polyhedron of dimension that equals the Krull dimension of M/P . Since M/P has prime characteristic and Krull dimension at most one, $\Sigma_{M/P}^c(Q)$ is the image of the projection to S^{n-1} of finitely many rays in \mathbb{R}^n and hence $\Sigma_{M/P}^c(Q)$ and $\Sigma_M^c(Q)$ are finite sets. Then there exists a cyclic subgroup Q' of Q such that $S(Q, Q') := \{\chi / \sim \mid \chi: Q \rightarrow \mathbb{R}, \chi \neq 0, \chi(Q') = 0\}$ is contained in $\Sigma_M^c(Q)$ and, by [6, Corollary 4.5], M is finitely generated as a $\mathbb{Z}Q'$ -module. We take q_0 to be a generator of Q' and define y_1, \dots, y_r to be generators of M as $\mathbb{Z}Q'$ -module. Note that according to [5] $\Delta_A(Q) = \Delta_M(Q)$ and hence A is m -tame if and only if M is m -tame.

Our aim from now on is to construct a finite set of characters V and to choose the negative integer β from the definition of the complex Y such that:

- (i) Y/G is a compact space;
- (ii) Y is $(m - 1)$ -acyclic;
- (iii) the stabilizer in G of a cell in Y is always polycyclic and hence of type FP_∞ .

The construction of Y resembles that of Åberg's complex in [1]; in fact Lemma 2.1(ii) allows us to use Åberg's argument [1, Chap. 3, Proposition 3.3] verbatim to show that if the set of characters V is $1, 2, \dots, (m-1)$ -domesticated then Y is $(m-1)$ -acyclic. We will construct the set V in such a way that all characters in V lie in $\Delta_M(Q)$ and by [1, Chap. 1, Proposition 6.3] if M is m -tame the set V is $1, 2, \dots, (m-1)$ -domesticated. This shows that Y is really $(m-1)$ -acyclic.

We need to explain the structure of Y and X as CW-complexes. If the set V contains d characters then X is built by gluing d -cubes of the type $\{\prod_{v \in V} [(a_v, r_v)] : z_v \leq r_v \leq z_v + 1\}$ on some parts of their faces, where a_v are elements of A and z_v are rational integers. These d -cubes form the d -cells in X and every i -cell ($i \leq d$) of X is an i -cube of the face of a d -cell in X . The intersection of Y with an i -cell in X contained in the d -cube $\{\prod_{v \in V} [(a_v, r_v)] : z_v \leq r_v \leq z_v + 1\}$ is homeomorphic to the intersection of the subspace $\text{im } \chi$ of $\prod_v \mathbb{R}_v$ with an i -cube contained in $\{\prod_v r_v \in \prod_v \mathbb{R}_v : z_v \leq r_v \leq z_v + 1\}$ and so is empty or homeomorphic to the unit j -cube B^j for some $j \leq i$. Now we define a j -cell of Y to be every intersection of Y with a cell in X which is homeomorphic to B^j .

2.3. The Construction of the Set of Characters V

Let w_i ($i = 1, 2$) be a character of Q' given by $w_i(q_0) = (-1)^{i+1}$ and let W_i ($i = 1, 2$) be the set of all characters v of Q extending w_i such that M is not finitely generated as $\mathbb{Z}Q_v$ -module. The last condition is equivalent to the graph X_v not being a single line and also equivalent to $v \in \Delta_M(Q) = \bigcup_j \Delta_{M/P_j}(Q)$ for a complete set of minimal prime ideals P_1, \dots, P_s in M . We claim that all the elements of W_i are discrete characters and W_i is a finite set for $i = 1, 2$.

Let $v \in \Delta_{M/P}(Q) \cap W_i$ for some minimal prime ideal P in M . Since M is finitely generated as $\mathbb{Z}[q_0^{\pm 1}]$ -module, for any element q of Q there is a relation $f_0(q_0)q^\alpha + f_1(q_0)q^{\alpha-1} + \dots + f_\alpha(q_0) = 0$ in M/P for some positive integer α and elements $f_i(q_0)$ of $\mathbb{Z}_p[q_0^{\pm 1}]$, where p is the characteristic of the ring M/P and \mathbb{Z}_p denotes the finite field consisting of p elements. As shown in [4, Lemma 3.2] there is a substitution $q_0 = aq^s$ for an integer s such that the above relation becomes of the type $g_0(a)q^\beta + g_1(a)q^{\beta-1} + \dots + g_\beta(a) = 0$ for some positive integer β and monomials $g_i(a) = z_i a^{\beta_i}$, $z_i \in \mathbb{Z}_p$, $0 \leq i \leq \beta$. Since M/P is not finitely generated as $\mathbb{Z}Q_v$ -module, by [5, Proposition 2.1] $v(a^{\beta_i} q^{\beta-i}) = v(a^{\beta_j} q^{\beta-j})$ for some $i < j$ such that the coefficients z_i and z_j are non-trivial. Then $v(q) = (\beta_j - \beta_i)v(q_0)/((j-i) + s(\beta_j - \beta_i)) \in \mathbb{Q} \subset \mathbb{R}$; note that the denominator cannot be trivial otherwise $\beta_j - \beta_i = 0$ and hence $i = j$, a contradiction. Therefore $v(q) \in \mathbb{Q}$ for every element q of Q , in particular for

elements of any finite generating set of Q . Thus the set W_i contains only discrete characters and is finite for each $i = 1, 2$.

Denote by V_i , for $i = 1, 2$, the set $\{d_v v : v \in W_i, d_v \text{ is a positive real number such that the image of } d_v v \text{ is exactly } \mathbb{Z}\}$ and let $V = V_1 \cup V_2$. We note that $\Sigma_A^c(Q) = \Sigma_M^c(Q) = V/\sim$.

2.4. G Acts Cocompactly on Y

A typical point of Y is $\prod_{v \in V} [(a_v, r_v)]$ where the a_v are elements of A and $\prod_v r_v \in \text{Im } \chi \subseteq \prod_v \mathbb{R}_v$. In Theorem 2.4 we shall prove that for any such point there exists an element a of A such that $\prod_v [(a_v, r_v)] = \prod_v [(a, r_v)]$. Since $(-a) * \prod_v [(a, r_v)] = \prod_v [(0, r_v)]$ we have $Y/A \simeq \text{Im } \chi$ and so $Y/G \simeq \text{Im } \chi/Q$, where Q acts on $\prod_v \mathbb{R}_v$ via the map χ and $Q \simeq \mathbb{Z}^n$ acts additively on \mathbb{R}^n . The map from $\mathbb{R}^n/\mathbb{Z}^n \simeq (Q \otimes \mathbb{R})/Q$ to $\text{Im } \chi/Q$ induced by χ is surjective and continuous and hence the space $\text{Im } \chi/Q \simeq Y/G$, being an image of a compact space, is also compact.

DEFINITION. If B is a subset of $\prod_v \mathbb{R}_v$ we define $[[B]] = \{\prod_v c_v \in \prod_v \mathbb{R}_v : \text{there exists } b = \prod_v b_v \in B \text{ such that } c_v - b_v \in [0, 1) \text{ and } c_v \in \mathbb{Z} \text{ for all } v \in V\}$.

Now we consider the case $B = \text{Im } \chi$.

LEMMA 2.3. $[[\text{Im } \chi]]$ is finite modulo the Q -action.

Proof. Let $\prod_v b_v \in \text{Im } \chi$, $[[\prod_v b_v]] = \prod_v c_v$, and let q be an element of Q . Then $[[q * \prod_v b_v]] = [[\prod_v (v(q) + b_v)]] = q * \prod_v c_v$ because $\prod_v v(q)$ is an integral point of $\prod_v \mathbb{R}_v$ and hence $[[\text{Im } \chi]]$ is closed under the Q -action.

Let d be the number of the valuations in V . Since $[[\text{Im } \chi]]$ consists of integral points $\prod_{v \in V} c_v$ and for every such point there exists an element $\prod_{v \in V} b_v$ of $\text{Im } \chi$ such that $0 \leq c_v - b_v < 1$ we have

$$[[\text{Im } \chi]] \subseteq \bigcup_{r \in \Delta} ((r + \text{Im } \chi) \cap \mathbb{Z}^d),$$

where $\Delta = \{r \in (\text{Im } \chi)^\perp : |r| \leq \sqrt{d}\}$. Since for every r of $\prod_v \mathbb{R}_v = \mathbb{R}^d$ the space $(r + \text{Im } \chi)/Q$ is compact and \mathbb{Z}^d/Q forms a discrete subset in \mathbb{R}^d/Q we have $((r + \text{Im } \chi) \cap \mathbb{Z}^d)/Q$ is finite. In order to prove Lemma 2.3 it is sufficient to show that the set $D = \{r \in \Delta : (r + \text{Im } \chi) \cap \mathbb{Z}^d \neq \emptyset\}$ is finite. After an \mathbb{R} -linear transformation of the coordinate system we can assume that

$$\begin{aligned} \text{Im } \chi &= \underbrace{\mathbb{R} \times \cdots \times \mathbb{R}}_{t \text{ times}} \times \mathbf{0} \times \cdots \times \mathbf{0}, \\ (\text{Im } \chi)^\perp &= \mathbf{0} \times \cdots \times \mathbf{0} \times \underbrace{\mathbb{R} \times \cdots \times \mathbb{R}}_{d-t \text{ times}} \end{aligned}$$

for some positive integer t . Denote by L the image of the integral points in \mathbb{R}^d under that transformation. Then L is a lattice in \mathbb{R}^d and

$$D \subseteq \{r \in (\operatorname{Im} \chi)^\perp : |r| \leq c, (r + \operatorname{Im} \chi) \cap L \neq \emptyset\}$$

for some constant c . Since all the characters of V are discrete, L has a \mathbb{Z} -basis of rational vectors and so $L \subseteq (\mathbb{Z}/b)^d$ for some positive integer b . Then D is contained in the discrete set

$$\begin{aligned} \{r \in (\operatorname{Im} \chi)^\perp : (r + \operatorname{Im} \chi) \cap (\mathbb{Z}/b)^d \neq \emptyset\} \\ = \mathbf{0} \times \cdots \times \mathbf{0} \times \underbrace{\mathbb{Z}/b \times \cdots \times \mathbb{Z}/b}_{d-t \text{ times}}. \end{aligned}$$

Since D is bounded we obtain that D is finite as required.

THEOREM 2.4. *For every $\prod_v s_v$ in $[[\operatorname{Im} \chi]]$ and every set $\{a_v \in A : v \in V\}$ there exists an element a in A such that $\prod_v [(a_v, s_v)] = \prod_v [(a, s_v)]$. Consequently for every $\prod_v r_v \in \operatorname{Im} \chi$ such that $[[\prod_v r_v]] = \prod_v s_v$ one has $\prod_v [(a_v, r_v)] = \prod_v [(a, r_v)]$.*

In order to prove Theorem 2.4 we need the following proposition.

PROPOSITION 2.5. *Let B be a finitely generated $\mathbb{Z}Q$ -module generated by the elements b_1, \dots, b_k with annihilator I in $\mathbb{Z}Q$ such that $\mathbb{Z}Q/I$ is a ring of non-zero characteristic and Krull dimension one. Let $\{v_1, \dots, v_r\}$ be a set of discrete characters of Q lying in an open half subspace of $\mathbb{R}^n = \operatorname{Hom}_{\mathbb{Z}}(Q, \mathbb{R})$ with all $\mathbb{R}^+ v_i$ representing different rays in \mathbb{R}^n . Let B_{v_i} be the $\mathbb{Z}Q_{v_i}$ -submodule of B generated by b_1, \dots, b_k and let q_{v_i} be an element of Q such that $v_i(q_{v_i}) = 1$ for $1 \leq i \leq r$. Then for any set $\{b_{v_i} : 1 \leq i \leq r\}$ of elements of B and any set of integers $\{s_{v_i} : 1 \leq i \leq r\}$ there exists an element b in B such that $b - b_{v_i} \in q_{v_i}^{s_{v_i}} B_{v_i}$ for all i .*

Deduction of Theorem 2.4 from Proposition 2.5. By Lemma 2.3 $[[\operatorname{Im} \chi]]$ is finite modulo the action of Q . Let $\prod_v s_v = T_i + \prod_v v(q)$ for an element q of Q and a point $T_i = \prod_v r_{v,i} \in \prod_v \mathbb{R}_v$ from a fixed representative set T_1, \dots, T_m of $[[\operatorname{Im} \chi]]$ modulo Q . We have

$$\prod_v [(a_v, s_v)] = \prod_v [(a, s_v)] \quad \text{for some } a \in A$$

if and only if

$$a_v \pi(q_v)^{s_v} G_v = a \pi(q_v)^{s_v} G_v \quad \text{for each } v \in V,$$

and this, in turn, is equivalent to

$$q_v^{-s_v} (a - a_v) \in G_v \cap A = q_v^\beta A_v \quad \text{for each } v \in V.$$

Since $s_v + \beta = v(q) + r_{v,i} + \beta \leq v(q) + \beta_1$ where $\beta_1 = \max\{r_{v,i} : 1 \leq i \leq m, v \in V\} + \beta$, it is sufficient to find $a' \in A$ such that $a' - a'_v \in q_v^{\beta_1} A_v$ for every $v \in V$ where $a'_v = q^{-1} a_v \in A$. Then $a = qa'$ is an element with the required properties.

Since $A = Ma_1 + \dots + Ma_s$ where $M = \mathbb{Z}Q/I$ and $I = \text{ann}_{\mathbb{Z}Q} A$, it suffices to consider the case $s = 1$ and $a'_v = 0$ for $v \neq w$ and $a'_w = ma_1$ for some element m of M and a character w of V . Thus we reduce the problem to finding m' in M such that $m' \in q_v^{\beta_1} \mathbb{Z}Q_v + I/I$ for $v \neq w$ and $m' - m \in q_w^{\beta_1} \mathbb{Z}Q_w + I/I$, and then $a' = m'a_1$ is an element with the required properties.

Without loss of generality we can assume that w is a character from V_1 (see the end of Section 2.3 for the definition of V_1) and by Proposition 2.5 there exists an element $x \in M$ such that $x - m \in q_w^{\beta_1} \mathbb{Z}Q_w + I/I$ and $x \in q_v^{\beta_1} \mathbb{Z}Q_v + I/I$ for all $v \neq w$ in V_1 . As shown in Section 2.2 we have that M is generated by y_1, \dots, y_r as $\mathbb{Z}[q_0^{\pm 1}]$ -module. From the very beginning we can assume that β is a sufficiently small negative integer such that $y_i \in q_v^{\beta_1} \mathbb{Z}Q_v + I/I$ for all $v \in V$ and $1 \leq i \leq r$. Then $x = \sum_{i=1}^r (f_i + g_i) y_i$ for some $f_i \in \mathbb{Z}[q_0]$ and $g_i \in \mathbb{Z}[q_0^{-1}]$. We take $m' = x - \sum_i f_i y_i = \sum_i g_i y_i$ and hence $m' - x \in q_v^{\beta_1} \mathbb{Z}Q_v + I/I$ for $v \in V_1$ and $m' \in q_v^{\beta_1} \mathbb{Z}Q_v + I/I$ for $v \in V_2$. By the choice of x we obtain that m' has the desired property.

Proof of Proposition 2.5. 1. Let I_1, \dots, I_k be ideals of $\mathbb{Z}Q$ containing I and suppose that the proposition holds whenever $I_i \subseteq \text{ann}_{\mathbb{Z}Q} B$. We show first that if there exists a chain

$$B = B_0 \supseteq B_1 \supseteq B_2 \supseteq \dots \supseteq B_m \supseteq B_{m+1} = 0$$

of $\mathbb{Z}Q$ -modules such that $\text{ann}_{\mathbb{Z}Q}(B_i/B_{i+1}) \supseteq I_{s(i)}$ for every i and some integers $1 \leq s(i) \leq k$ then the proposition holds.

Let $\{b_{i,1}, \dots, b_{i,m_i}\}$ be a set of generators of B_i as $\mathbb{Z}Q$ -module. For a sufficiently large positive integer u we have $q_0^u b_{i,j} \in B_{v_t}$ for any $1 \leq t \leq r$, $1 \leq j \leq m_i$, $0 \leq i \leq m$, where q_0 is an element of Q such that $v_t(q_0) > 0$ for all t . Since the proposition holds for B_i/B_{i+1} generated by $\{q_0^u b_{i,j} + B_{i+1} : 1 \leq j \leq m_i\}$ as $\mathbb{Z}Q$ -module we have that for any set of integers $\{s_t : 1 \leq t \leq r\}$ and any set $\{a_{i,t} : 1 \leq t \leq r\}$ of elements of B_i there exists an element λ_i in B_i such that

$$\lambda_i - a_{i,t} \in q_{v_t}^{s_t} \left(\sum_{j=1}^{m_i} \mathbb{Z}Q_{v_t} q_0^u b_{i,j} \right) + B_{i+1} \subseteq q_{v_t}^{s_t} B_{v_t} + B_{i+1}.$$

Using this property we find elements c_i in B_i such that $c_0 - b_{v_t} = d_{t,1} + c_{t,1}$, $c_i + c_{t,i} = d_{t,i+1} + c_{t,i+1}$ where $c_{t,i} \in B_i$, $d_{t,i} \in q_{v_t}^{s_t} B_{v_t}$ for $1 \leq i \leq m$, $1 \leq t \leq r$. Since $B_{m+1} = 0$ we have $c_{t,m+1} = 0$ and hence $b := c_0 + c_1 +$

$\cdots + c_m = b_{v_i} + d_{t,1} + \cdots + d_{t,m+1} \in b_{v_i} + q_{v_i}^{s_t} B_{v_i}$ for all t and therefore b has the required property.

2. In general $I \subseteq \sqrt{I} = P_1 \cap \cdots \cap P_s$ for some prime ideals P_1, \dots, P_s in $\mathbb{Z}Q$ such that the Krull dimension of the rings $\mathbb{Z}Q/P_i$ is less than or equal to one. Let $\sqrt{I}^{m+1} \subseteq I$. Then

$$B \supseteq \sqrt{I}B \supseteq \sqrt{I}^2 B \supseteq \cdots \supseteq \sqrt{I}^{m+1} B = 0$$

and for the $\mathbb{Z}Q$ -modules $B_i = \sqrt{I}^i B / \sqrt{I}^{i+1} B$ the following inclusions hold

$$B_i \supseteq P_1 B_i \supseteq (P_1 P_2) B_i \supseteq \cdots \supseteq (P_1 \cdots P_s) B_i = 0.$$

According to step 1 it is sufficient to consider the case $P \subseteq \text{ann}_{\mathbb{Z}Q} B$ for some prime ideal P of $\mathbb{Z}Q$ such that the Krull dimension of $\mathbb{Z}Q/P$ is less than or equal to one.

In the case when the Krull dimension of $\mathbb{Z}Q/P$ is zero the ring $\mathbb{Z}Q/P$ is finite and hence B is finite. Then B is finitely generated as $\mathbb{Z}Q_v$ -module and by [5, Proposition 2.1], $B = B_v = q_{v_i}^{s_v} B_v$ for any integer s_v and any real character v of Q . Then every b in B has the required property.

3. It remains to consider the case $P \subseteq \text{ann}_{\mathbb{Z}Q} B$ where $\mathbb{Z}Q/P$ is an integral domain of Krull dimension one and non-zero characteristic. Then B is p -torsion for some prime integer p and $B = \mathbb{Z}_p Q b_1 + \cdots + \mathbb{Z}_p Q b_s$. With abuse of notation we write P for the image of the ideal P in $\mathbb{Z}_p Q$. To finish the proof of the proposition it suffices to consider the case $s = 1$ and $b_{v_i} = 0$ for $i \neq j$ and $b_{v_j} \in Q$ for some j between 1 and r . We aim to find an element λ in $\mathbb{Z}_p Q$ such that $\lambda \in q_{v_i}^{s_{v_i}} \mathbb{Z}_p Q_{v_i} + P$ for $i \neq j$ and $\lambda - b_{v_j} \in q_{v_j}^{s_{v_j}} \mathbb{Z}_p Q_{v_j} + P$. Now multiplying by $b_{v_j}^{-1}$ and changing the integers s_{v_i} to $s'_{v_i} = s_{v_i} - v_i(b_{v_j})$ we have to consider the above inclusions with $b_{v_j} = 1_Q$ and the new set of integers s'_{v_i} . The following lemma finishes off the proof of Proposition 2.5.

LEMMA 2.6. *In the previous notation for any set of integers $\{s_{v_i}; 1 \leq i \leq r\}$ there exists an element λ of $\mathbb{Z}_p Q$ such that $\lambda \in q_{v_i}^{s_{v_i}} \mathbb{Z}_p Q_{v_i} + P$ for $1 \leq i \neq j \leq r$ and $\lambda - 1_Q \in q_{v_j}^{s_{v_j}} \mathbb{Z}_p Q_{v_j} + P$.*

Proof. Since $\text{Hom}_{\mathbb{Z}}(Q, \mathbb{Z}) \simeq \mathbb{Z}^n \simeq Q \subset \mathbb{R}^n$ we can consider the discrete characters v_1, \dots, v_r as elements of the group $Q \subset \mathbb{R}^n$ and v_i as maps given by $v_i(q) = (v_i, q)$, where $(,)$ is the standard inner product in \mathbb{R}^n . We prove Lemma 2.6 by induction on the rank of Q and assume for the moment that Lemma 2.6 holds for $r(Q) \leq 2$. Let $n \geq 3$ be the rank of Q . We want to find a hyperplane L in $\mathbb{R}^n = Q \otimes \mathbb{R}$ defined by $(h, L) = 0$ for some rational vector h in \mathbb{R}^n such that

(1) if $\pi: \mathbb{R}^n \rightarrow L$ is the projection map then all projections $\pi(v_i)$ are non-trivial;

(2) the rays in $\mathbb{R}^{n-1} \simeq L$ determined by $\pi(v_i)$ are all different and lie in an open half subspace of L .

Note that $\mathbb{R}\pi(v_i) = \mathbb{R}\pi(v_j)$ if and only if h, v_i, v_j are linearly dependent.

Let $h_1 = q_0 + \epsilon e$ where e is a unit vector in \mathbb{R}^n , ϵ a real number, and q_0 is an element of Q such that $v_i(q_0) > 0$ for all i . If $|\epsilon|$ is sufficiently small then $(v_i, h_1) > 0$ for all i . If in addition $(h, h_1) = 0$ then $(\pi(v_i), h_1) = (v_i, h_1) > 0$ for all i , and so all $\pi(v_i)$ lie in the open half subspace $\{l \in L: (l, h_1) > 0\}$ of L . Thus if $h \in \bigcup_{|h_1 - q_0| \leq \epsilon} h_1^\perp \setminus \bigcup_{i \neq j} \text{span}\{v_i, v_j\}$ and h is a rational vector in \mathbb{R}^n , the hyperplane L defined by h has the required properties. Since the dimension of any of the spaces h_1^\perp is $n - 1$, and thus at least 2, and the first union is infinite we have that $\bigcup_{|h_1 - q_0| \leq \epsilon} h_1^\perp \setminus \bigcup_{i \neq j} \text{span}\{v_i, v_j\}$ is non-empty and obviously the rational points are dense in it. Finally we obtain a hyperplane L with the required properties.

Now let w_i be integral vectors lying on the rays defined by $\pi(v_i)$ and we consider w_i as discrete characters of $Q_1 = L \cap \mathbb{Z}^n \simeq \mathbb{Z}^{n-1} \subset Q \simeq \mathbb{Z}^n$ defined by $w_i(q) = (w_i, q)$ for $q \in Q_1$. We apply induction for the group Q_1 . Then there exists an element λ in $\mathbb{Z}_p Q_1 \subseteq \mathbb{Z}_p Q$ such that $\lambda \in q_{w_i}^{\alpha_i} \mathbb{Z}_p(Q_1)_{w_i} + (P \cap \mathbb{Z}_p Q_1)$ for $i \neq j$ and $\lambda - 1 \in q_{w_j}^{\alpha_j} \mathbb{Z}_p(Q_1)_{w_j} + (P \cap \mathbb{Z}_p Q_1)$ where q_{w_i} are elements of Q_1 such that $w_i(q_{w_i}) = 1$, α_i are rational integers, and $(Q_1)_{w_i} = \{q \in Q_1 \mid w_i(q) \geq 0\}$. Since $v_i(q_{w_i}) > 0$, we have $q_{w_i}^{\alpha_i} \mathbb{Z}_p(Q_1)_{w_i} \subseteq q_{v_i}^{s_{v_i}} \mathbb{Z}_p Q_{v_i}$ for a sufficiently big α_i and then λ is an element with the required properties.

Now we consider the case $n = 2$ (if $n = 1$ then r should be 1 as well so there is nothing to prove). We write x, y for the standard generators of $Q = \mathbb{Z}^2 \subset \mathbb{R}^2$ and without loss of generality we can assume that $v_i(y) > 0$ for all i . By reordering the set $\{v_1, v_2, v_3, \dots, v_{r-1}, v_r\}$ if necessary we can assume that $v_i(x)/v_i(y) < v_{i+1}(x)/v_{i+1}(y)$ for $1 \leq i \leq r - 1$. Since $\mathbb{Z}_p Q_{v_i} \not\equiv \mathbb{Z}_p Q \pmod{P}$ if and only if v_i belongs to $\Delta_{\mathbb{Z}_p Q/P}(Q)$ (see [5, Proposition 2.1]) we can assume without loss of generality that all v_i have this property. By enlarging the set of characters $\{v_1, \dots, v_r\}$ we can assume as well that the classes of v_1, \dots, v_r represent all the elements in $\Sigma_{\mathbb{Z}_p Q/P}^c(Q)$ lying in the open half subspace of \mathbb{R}^2 defined by y .

Since the Krull dimension of $\mathbb{Z}_p Q/P$ equals one there exists a polynomial $t \in \mathbb{Z}_p[x, y]$ such that $P = (t)$. Let M be the convex hull of the support of t in $\mathbb{R}^2 = \mathbb{R} \otimes_{\mathbb{Z}} Q$. We order clockwise the edges of the boundary of M consecutively l_1, \dots, l_s and by [6] for every i between 1 and r there exists $\sigma(i)$ such that v_i is perpendicular to $l_{\sigma(i)}$ and for any $m \in M \cap Q$ and $q \in l_{\sigma(i)} \cap Q$ we have $v_i(m) \geq v_i(q)$. Using the description of the geometrical invariant $\Sigma_c(Q)$ in [6] for one related kQ -module

C where k is a commutative ring with 1, the field \mathbb{Z}_p in our case, and reordering l_1, \dots, l_s by translating the indices modulo s if necessary, we can assume $\sigma(i) = i$ for $i \leq r$.

Let P_i be the intersection of the edges l_{i-1} and l_i in \mathbb{R}^2 where $l_0 = l_s$ and let q_1, \dots, q_s be the elements of $Q \simeq \mathbb{Z}^2$ corresponding to the points P_1, \dots, P_s . Let $t_1 = c_1^{-1} q_j^{-1} t$ and $t_2 = c_2^{-1} q_{j+1}^{-1} t$ where each c_i is the coefficient of q_{j+i-1} in t . If v is a vector in \mathbb{R}^2 and $g = \sum_q z_q q$ is an element in $\mathbb{Z}_p Q$ we define $g_v = \sum_{(q,v) \geq 0} z_q q$ and $g_v^+ = \sum_{(q,v) > 0} z_q q$. Denote $\lambda_1 = (t_2)_{x^{-1}} + (t_1)_x - 1_Q$. Now $\lambda_1 \equiv ((t_2)_{x^{-1}} - 1_Q) - (t_1)_{x^{-1}}^+ \pmod{P}$ and the elements of the supports of $(t_1)_{x^{-1}}^+$ and $(t_2)_{x^{-1}} - 1_Q$ lie in the open half subspaces defined by v_i for $1 \leq i \leq j-1$. We have as well $\lambda_1 \equiv ((t_1)_x - 1_Q) - (t_2)_x^+ \pmod{P}$ and the elements of the supports of $(t_1)_x - 1_Q$ and $(t_2)_x^+$ lie in the open half subspaces of \mathbb{R}^2 defined by v_i for $j+1 \leq i \leq r$. Thus $\lambda_1 \in \mathbb{Z}_p Q_{v_i}^+ + P$ for $1 \leq i \neq j \leq r$ where $Q_{v_i}^+ = \{q \in Q : v_i(q) > 0\}$.

By the construction of λ_1 , the support of $\lambda_1 - 1_Q$ lies in the open half subspace of \mathbb{R}^2 defined by v_j so $\lambda_1 - 1_Q \in \mathbb{Z}_p Q_{v_j}^+ + P$. Since $(\lambda_1 - 1_Q)^{p^k} = \lambda_1^{p^k} - 1_Q$, for a sufficiently large positive integer k we obtain that $\lambda = \lambda_1^{p^k}$ has the required properties. This finishes the proof of Lemma 2.6.

As a corollary of Theorem 2.4 we get

COROLLARY 2.7. *Y/G is a compact space.*

2.5. Stabilizers in G of Cells in Y

We aim to show that if P is the stabilizer of a vertex $\prod_{v \in V} g_v G_v$ in X lying in $f^{-1}([\text{im } \chi])$ then P is polycyclic and hence of type FP_∞ . As we will see later the stabilizer in G of a cell in Y is always of the form described above.

PROPOSITION 2.8. *The intersection $P \cap A$ is a finite group.*

Proof. If a is an element of A stabilizing a vertex $g_v G_v$ of X_v where $g_v = b\pi(q_v)^z$ for some element b of A and some rational integer z , we have $(b\pi(q_v)^z)^{-1} a b\pi(q_v)^z = q_v^{-z} a \in G_v \cap A = q_v^\beta A_v$ and so $a \in q_v^{\beta + \alpha_v(g_v)} A_v$. Then $P \cap A \subseteq \bigcap_{v \in V} q_v^\alpha A_v$ for some sufficiently small integer α .

We claim that if B_v are finitely generated $\mathbb{Z}Q_v$ -submodules of A then the intersection $\bigcap_{v \in V} B_v$ is always finite. Then Proposition 2.8 obviously holds. We divide the proof into several steps.

(1) Let $I = \text{ann}_{\mathbb{Z}Q} A$ and let $I_0 \subseteq I_1 \subseteq \dots \subseteq I_j = \mathbb{Z}Q$ be an increasing sequence of ideals in $\mathbb{Z}Q$ such that $I_0 \subseteq I$. Then $((\cap_v B_v) \cap I_i A) / ((\cap_v B_v) \cap I_{i-1} A)$ embeds in $\cap_v ((B_v \cap I_i A) + I_{i-1} A / I_{i-1} A) \subseteq I_i A / I_{i-1} A$. We assume the claim holds for the $\mathbb{Z}Q$ -modules $I_i A / I_{i-1} A$ for $1 \leq i \leq j$. Since the ring $\mathbb{Z}Q_v$ is Noetherian for a discrete character v , we have that $B_v \cap I_i A$ is finitely generated as $\mathbb{Z}Q_v$ -module and therefore the intersection $\cap_v ((B_v \cap I_i A) + I_{i-1} A / I_{i-1} A)$ is finite for $1 \leq i \leq j$. Then all the quotients $((\cap_v B_v) \cap I_i A) / ((\cap_v B_v) \cap I_{i-1} A)$ for $1 \leq i \leq j$ are finite and then $\cap_v B_v$ is finite as required.

(2) Let the radical \sqrt{I} of I be the intersection of some prime ideals P_1, \dots, P_s in $\mathbb{Z}Q$ and $\sqrt{I}^k \subseteq I$ for some positive integer k . Then the Krull dimension of the rings $\mathbb{Z}Q/P_i$ for $1 \leq i \leq s$ is less than or equal to 1. The sequence of ideals in $\mathbb{Z}Q$, $\sqrt{I}^k \subseteq \sqrt{I}^{k-1} \subseteq \dots \subseteq \sqrt{I} \subseteq \mathbb{Z}Q$, and the above remark allow us to assume that $P_1 \cap \dots \cap P_s \subseteq \text{ann}_{\mathbb{Z}Q} A$. In this case the following sequence of ideals in $\mathbb{Z}Q$, $P_1 \cap \dots \cap P_s \subseteq P_1 \cap \dots \cap P_{s-1} \subseteq \dots \subseteq P_1 \subseteq \mathbb{Z}Q$, shows that we can further assume that $\text{ann}_{\mathbb{Z}Q} A$ contains a prime ideal P of $\mathbb{Z}Q$ such that the Krull dimension of $\mathbb{Z}Q/P$ is less than or equal to one.

(3) Let $A = \mathbb{Z}Qa_1 + \dots + \mathbb{Z}Qa_s$ and $P \subseteq \text{ann}_{\mathbb{Z}Q} A$ as above. We use induction on s to show that if B_v are finitely generated $\mathbb{Z}Q_v$ -submodules of A then the intersection $\cap_{v \in V} B_v$ is finite.

(3.1) We assume $s > 1$ and the statement is true for $\mathbb{Z}Q/P$ -modules generated by at most $s-1$ elements. Let $J = \{\lambda \in \mathbb{Z}Q : \lambda a_1 \in \mathbb{Z}Qa_2 + \dots + \mathbb{Z}Qa_s\}$. If P is strictly contained in J then either the radical \sqrt{J} of the ideal J is an intersection of finitely many prime ideals Q_1, \dots, Q_t in $\mathbb{Z}Q$ strictly containing P or $\sqrt{J} = \mathbb{Z}Q$. In the first case Q_1, \dots, Q_t are maximal ideals of $\mathbb{Z}Q$ and hence they are cofinite in $\mathbb{Z}Q$. In both cases $\mathbb{Z}Q/J$ is a finite ring and so $A/\mathbb{Z}Qa_2 + \dots + \mathbb{Z}Qa_s$ is finite. Then $(\cap_{v \in V} B_v) / ((\mathbb{Z}Qa_2 + \dots + \mathbb{Z}Qa_s) \cap (\cap_v B_v))$ is finite and by induction $(\cap_{v \in V} B_v) \cap (\mathbb{Z}Qa_2 + \dots + \mathbb{Z}Qa_s)$ is finite again. We obtain that $\cap_{v \in V} B_v$ is finite as required.

In the case when $P = J$, $A = \mathbb{Z}Qa_1 \oplus (\mathbb{Z}Qa_2 + \dots + \mathbb{Z}Qa_s)$. Let $\{b_1, b_2, \dots, b_i\}$ be a finite set containing a generating set of B_v as $\mathbb{Z}Q_v$ -module for all $v \in V$, so $B_v \subseteq \sum_{i=1}^t \mathbb{Z}Q_v b_i$. We write $b_i = b_{i,1} + b_{i,2}$ where $b_{i,1} \in \mathbb{Z}Qa_1$ and $b_{i,2} \in \mathbb{Z}Qa_2 + \dots + \mathbb{Z}Qa_s$. Thus $\cap_v B_v \subseteq (\cap_v B_{v,1}) \oplus (\cap_v B_{v,2})$ where $B_{v,j} = \sum_i \mathbb{Z}Q_v b_{i,j}$ for $j = 1, 2$ and $v \in V$. By induction both $\cap_v B_{v,1}$ and $\cap_v B_{v,2}$ are finite and hence $\cap_v B_v$ is finite.

(3.2) It remains to consider the case $s = 1$, $A \simeq \mathbb{Z}Q/I$ for an ideal I in $\mathbb{Z}Q$ containing P where $\mathbb{Z}Q/P$ is an integral domain of non-zero characteristic and Krull dimension 1. If $P \neq I$ then A is finite and there is nothing to prove. If $I = P$ then $B_v \subseteq q_v^r \mathbb{Z}Q_v + I/I$ for a sufficiently small

negative integer r and every $v \in V$ where q_v is an element of Q such that $v(q_v) = 1$. We remind the reader that the ring $M = \mathbb{Z}Q/\text{ann}_{\mathbb{Z}Q} A$ for the initial $\mathbb{Z}Q$ -module A is finitely generated as $\mathbb{Z}[q_0^{\pm 1}]$ -module (see Sect. 2.2) and hence the new ring A is finite over the subring $S = \mathbb{Z}[q_0^{\pm 1}] + I/I$. Since A is infinite $S = \mathbb{Z}[q_0^{\pm 1}] + I/I$ is isomorphic to the Laurent polynomial ring $\mathbb{Z}_p[q_0^{\pm 1}]$, where p is the characteristic of the integral domain $A = \mathbb{Z}Q/I$.

Let V'_i ($i = 1, 2$) be the set of all valuations of the field of fractions of A with the group of rational integers \mathbb{Z} as value group with the additional property that the restriction of every element of V'_i to the Laurent polynomial ring $S \simeq \mathbb{Z}_p[q_0^{\pm 1}]$ is given by minimal formula and the value of q_0 is positive for $i = 1$ and negative for $i = 2$. A valuation v of $\mathbb{Z}_p[q_0^{\pm 1}]$ is given by minimal formula if $v(\sum_{k \in \mathbb{Z}} z_k q_0^k) = \min\{v(q_0^k): z_k \neq 0\}$ where $z_k \in \mathbb{Z}_p$. Then the restrictions of the valuations in V'_i to Q give some elements of V_i . We set $V' = V'_1 \cup V'_2$.

Now

$$\bigcap_{v \in V'} B_v \subseteq \{a \in A \mid v(a) \geq r, v \in V'\} =: C'.$$

Since $v(q_0) \geq 1$ for every $v \in V'_1$ we have

$$q_0^{-r} C' \subseteq C_1 := \{a \in A \mid v(a) \geq 0 \text{ if } v \in V'_1, v(a) \geq c \text{ if } v \in V'_2\}$$

for a negative integer $c \leq r(1 - \min\{v(q_0): v \in V'_2\})$. Thus $C_1 \subseteq \{a \in A: v(a) \geq 0, v \in V'_1\}$ and by [8, Chap. 6.1.3, Theorem 3] $\{a \in A: v(a) \geq 0, v \in V'_1\} = \{a \in A: a \text{ is integral over } O_{v_1}\}$ where $O_{v_1} = \{k \in \text{the field of fractions of } S: v_1(k) \geq 0\}$, v_1 is the valuation on the field of fractions of S given by minimal formula on S , and $v_1(q_0) = 1$. Since A is integral over $S = \mathbb{Z}_p[q_0^{\pm 1}]$ and O_{v_1} is a principal ideal domain, and so a unique factorization domain, any element of C_1 is integral over $S \cap O_{v_1} = \mathbb{Z}_p[q_0] =: E$. Thus $C_1 \subseteq D$ where D is the integral closure of E in the field of fractions K of A . By [8, Chap. 5.3.2, Theorem 2], D is a finitely generated E -module, say $D = Ed_1 + \cdots + Ed_m$. We define

$$D'_k = \left\{ e_1 d_1 + \cdots + e_m d_m \in D \mid e_i = \sum_{0 \leq \alpha < k} z_{\alpha, i} q_0^\alpha, z_{\alpha, i} \in \mathbb{Z}_p \right\},$$

$$D''_k = \left\{ e_1 d_1 + \cdots + e_m d_m \in D \mid e_i = \sum_{\alpha \geq k} z_{\alpha, i} q_0^\alpha, z_{\alpha, i} \in \mathbb{Z}_p \right\}.$$

We show that for sufficiently large k , both $C_1 \cap D'_k$ and $C_1/(C_1 \cap D''_k)$ are finitely generated \mathbb{Z}_p -modules and consequently $C_1, C', \bigcap_{v \in V'} B_v$, and $A \cap P$ are finite.

Generally D/D_k'' is a finitely generated \mathbb{Z}_p -module so we need to show that $C_1 \cap D_k''$ is finitely generated for sufficiently large k . Let $l \in C_1 \cap D_k''$ and so $l = q_0^k l_1$ for some $l_1 \in D$ and $v(l_1) = v(l) - kv(q_0) \geq c - kv(q_0) \geq 0$ for sufficiently large k and for all v in V_2' . Thus $C_1 \cap D_k'' \subseteq q_0^k \{m \in D: v(m) \geq 0, v \in V_2'\} = q_0^k \{\text{the integral closure of } \mathbb{Z}_p \text{ in } K\}$. The integral closure of \mathbb{Z}_p in K is finitely generated as \mathbb{Z}_p -module by [8, Chap. 5.3.2, Theorem 2] because the algebraic closure of \mathbb{Z}_p in K is a finite extension of \mathbb{Z}_p . This completes the proof of Proposition 2.8.

LEMMA 2.9. *If Γ is a cell of Y then the stabilizer of Γ in G coincides with the stabilizer in G of a point of $f^{-1}([\text{Im } \chi])$.*

Proof. Let Γ be the intersection of an i -subcell of the d -cell $J = \{\prod_v [(a_v, r_v)]: z_v \leq r_v \leq z_v + 1\}$ in X with Y , let $\gamma_1, \dots, \gamma_s$ be the vertices (0-subcells) of Γ , and let $g \in G$ stabilize the cell Γ . Then g permutes the vertices $\gamma_1, \dots, \gamma_s$, say $g * \gamma_i = \gamma_{\rho(i)}$ for some permutation ρ . If $\gamma_i = \prod_v [(a_v, s_{v,i})]$ we have $s_{v,i} + \alpha_v(g) = s_{v,\rho(i)}$ and so $\sum_i (s_{v,i} + \alpha_v(g)) = \sum_i s_{v,\rho(i)} = \sum_i s_{v,i}$. Hence $\alpha_v(g) = 0$ for $v \in V$ and $\gamma_{\rho(i)} = \gamma_i$ for all i . Thus g stabilizes the vertices $\gamma_1, \dots, \gamma_s$ and hence stabilizes Γ pointwise.

We note that the element g in G stabilizes the point $\prod_v [(a_v, r_v)]$ of X if and only if g stabilizes the vertex $\prod_v [(a_v, s_v)]$ of X where $\prod_v s_v = [\prod_v r_v]$, i.e., $s_v \in \mathbb{Z}$ and $s_v - r_v \in [0, 1)$ for all v . We obtain that $\{g \in G: g \text{ stabilizes } \Gamma \text{ pointwise}\} = \{g \in G: g \text{ stabilizes all points from the set } \Lambda\}$ where $\Lambda = \{\prod_v [(a_v, s_v)]: \text{there exists a point } \prod_v [(a_v, r_v)] \text{ in } \Gamma \text{ such that } \prod_v s_v = [\prod_v r_v]\} \subseteq \text{vertices in } X \text{ of the } d\text{-cell } J$. Let $\Lambda = \{T_1, \dots, T_r\}$ and $f(T_i) = [f(S_i)]$ for some points S_1, \dots, S_r of Γ and $T_j = \prod_v [(a_v, z_{v,j})]$ for $1 \leq j \leq r$. We claim that the point $T = \prod_v [(a_v, z_v)]$ where $z_v = \max\{z_{v,j}: 1 \leq j \leq r\}$ belongs to Λ . Indeed $(1/r)(S_1 + \dots + S_r)$ is a point of Γ and $[f((1/r)(S_1 + \dots + S_r))] = f(T)$. Thus $\{g \in G: g \text{ stabilizes } \Gamma \text{ pointwise}\} = \{g \in G: g \text{ stabilizes the vertex } T \text{ of } X\}$ and T is a point of $f^{-1}([\text{Im } \chi])$.

3. PROOF OF THEOREM B

3.1. Åberg's Construction

From now on we work in the notation of Theorem B. Without loss of generality we can assume Q is free abelian of rank n . To prove Theorem B we follow Åberg's construction from [1].

Let P_1, \dots, P_s be all the minimal associated primes for the $\mathbb{Z}Q$ -module A and so $P_i = \text{ann}_{\mathbb{Z}Q}(a_i)$ for some elements a_1, \dots, a_s of A . There is a $\mathbb{Z}Q$ -module homomorphism $i: N = \mathbb{Z}Q/P_1 \oplus \dots \oplus \mathbb{Z}Q/P_s \rightarrow A$ which sends $\lambda_1 \oplus \dots \oplus \lambda_s$ to $\lambda_1 a_1 + \dots + \lambda_s a_s$. By [1, Chap. 2, Proposition 1.4], i is an injection. From now on we consider N as a submodule of A .

We assume A is not m -tame but it is $(m-1)$ -tame, and, as in the introduction of [11], there exist m discrete characters $\chi_1, \chi_2, \dots, \chi_m$ of Q each with value group a subgroup of the group of rational integers \mathbb{Z} . Moreover for every $1 \leq i \leq m$ at least one of the rings $\mathbb{Z}Q/P_j$, $1 \leq j \leq s$, is not finitely generated as $\mathbb{Z}Q_{\chi_i}$ -module, where $Q_{\chi_i} = \{q \in Q: \chi_i(q) \geq 0\}$, and the convex hull of $\chi_1, \chi_2, \dots, \chi_m$ in $\text{Hom}_{\mathbb{Z}}(Q, \mathbb{R}) \simeq \mathbb{R}^n$ is an $(m-1)$ -dimensional simplex containing the origin as an internal point. We note that if z_1, z_2, \dots, z_m are positive integers then the characters $z_1 \chi_1, \dots, z_m \chi_m$ still have the described properties. By multiplying χ_m with a sufficiently big positive integer, if necessary, we can assume that χ_m is an integral combination of $\chi_1, \dots, \chi_{m-1}$.

We view Q as the integral lattice \mathbb{Z}^n in \mathbb{R}^n . Then there is an isomorphism $\theta: \text{Hom}_{\mathbb{Z}}(Q, \mathbb{Z}) \rightarrow \mathbb{Z}^n = Q$ given by $\chi(q) = (\theta(\chi), q)$ for all $q \in Q$ and each character χ of Q , where (\cdot, \cdot) is the standard inner product in \mathbb{R}^n . Denote the subgroup of Q generated by $\theta(\chi_1), \dots, \theta(\chi_m)$ by Q' and let Q'' be the subgroup of $Q = \mathbb{Z}^n$, consisting of all elements in Q perpendicular to Q' with respect to the standard inner product in \mathbb{R}^n . Then $Q' \times Q''$ is a subgroup of finite index in Q and since the FP_m -Conjecture holds for a group G if and only if it holds for a subgroup of finite index in G we can assume $Q = Q' \times Q''$.

Åberg notices in [1] that if G is of type FP_m the canonical map

$$H^0(Q', E) = E^{Q'} \rightarrow H_0(Q'', E) = E_{Q''}$$

is trivial where E stands for $(\mathbb{Z}G^{\mathbb{N}})_A$. We aim to construct an element of the Cartesian product $\mathbb{Z}G^{\mathbb{N}}$ whose images β in E is Q' -invariant and the image of β in $E_{Q''}$ is non-trivial. We need some properties of the elements of $\mathbb{Z}G^{\mathbb{N}}$ which vanish in $(\mathbb{Z}G^{\mathbb{N}})_{G''}$, where G'' is the subgroup of G containing A and $G''/A \simeq Q''$.

Let $\{F^d\}_{d \in \mathbb{Z}}$ be a decreasing filtration of A such that $\bigcup_{d \in \mathbb{Z}} F^d = A$. For an element f of the augmentation ideal of the group algebra $\mathbb{Z}A$ there is an order function $o(f)$ (see [1]) given by

$$o(f) = \sup\{d: f \equiv 0 \text{ in } \mathbb{Z}[A/F^d]\} \in \mathbb{Z} \cup +\infty.$$

The following proposition in the split case is proved in [1, 11].

PROPOSITION 3.1. *Let $A \rightarrow G \rightarrow Q$ be a short exact sequence of groups with A, Q abelian, G finitely generated, and let $\pi: Q \rightarrow G$ be a lifting of the projection map $G \rightarrow Q$. Let $\{F^d\}_{d \in \mathbb{Z}}$ be a decreasing filtration of $\mathbb{Z}Q''$ -submodules of A such that $\bigcup_{d \in \mathbb{Z}} F^d = A$ and let $\alpha = (\alpha_j)_{j \in \mathbb{N}}$ be an element of $\mathbb{Z}G^{\mathbb{N}}$ with $\alpha_j = \sum_q \alpha_{j,q} \pi(q)$, $\alpha_{j,q} \in \mathbb{Z}A$, $q \in Q'$. If α vanishes in $(\mathbb{Z}G^{\mathbb{N}})_{G''}$, where G'' is the subgroup of G generated by A and $\pi(Q'')$, then $\alpha_{j,q}$ are elements of the augmentation ideal of $\mathbb{Z}A$ and $\inf\{o(\alpha_{j,q}): j, q\} > -\infty$.*

Remark. In the case when all F^d are finitely generated $\mathbb{Z}Q''$ -modules and $\inf\{o(\alpha_{j,q}): j, k\} > -\infty$ it can be shown that α vanishes in $(\mathbb{Z}G^{\mathbb{N}})_{G''}$ (a split version is proved in [1, 11]) but as the filtration we will consider later consists of $\mathbb{Z}Q''$ -modules which are not finitely generated, we abstain from claiming sufficiency.

Proof of Proposition 3.1. We follow the proof of [11, Chap. 2.3, Proposition]. Let α vanish in $(\mathbb{Z}G^{\mathbb{N}})_{G''}$ and $\alpha = \sum_{i=1}^s (g_i'' - 1)\beta_i$ for some $g_i'' \in G''$, $\beta_i \in \mathbb{Z}G^{\mathbb{N}}$. Let $\beta_i = (\beta_{i,j})_{j \in \mathbb{N}}$ where $\beta_{i,j} = \sum_q \beta_{i,j,q} \pi(q)$ and $\beta_{i,j,q} \in \mathbb{Z}G''$, $q \in Q'$. Then $\alpha_{j,q} = \sum_{i=1}^s (g_i'' - 1)\beta_{i,j,q}$ for any positive number j and an element q of Q' . We write $g_i'' = a_i \pi(q_i'')$ for some $a_i \in A$ and $q_i'' \in Q''$, and define P to be the subgroup of G'' generated by $\pi(Q'')$ and $\{a_i: 1 \leq i \leq s\}$. Then $\alpha_{j,q}$ vanishes in $\mathbb{Z}[G''/P] \simeq \mathbb{Z}[A/(A \cap P)]$; here $\mathbb{Z}[G''/P]$ and $\mathbb{Z}[A/(A \cap P)]$ are the free abelian groups on the right coset classes of P in G'' and $A \cap P$ in A , respectively, and the isomorphism is of free abelian groups given by the obvious bijection between the coset classes. Since $P \cap A$ is finitely generated as $\mathbb{Z}Q''$ -module it is contained in some F^d and so $\alpha_{j,q}$ vanishes in $\mathbb{Z}[A/F^d]$. Thus $\alpha_{j,q}$ are elements of the augmentation ideal of $\mathbb{Z}A$ and $o(\alpha_{j,q}) \geq d$ for all j, q . Later we will need the following lemma.

LEMMA 3.2. *Let A be a $\mathbb{Z}Q$ -module, let R be a Noetherian subring of $\mathbb{Z}Q$, and let N be an R -submodule of A . If $\{F^d\}_{d \in \mathbb{Z}}$ is a decreasing filtration of R -submodules of N such that $\bigcup_d F^d = N$ then there exists a decreasing filtration $\{F_*^d\}_{d \in \mathbb{Z}}$ of R -submodules of A such that $F_*^d \cap N = F^d$ and $\bigcup_{d \in \mathbb{Z}} F_*^d = A$.*

Proof. Let

$$N = N_0 \subseteq N_1 \subseteq \cdots \subseteq N_\lambda \subseteq \cdots \subseteq N_\beta = A$$

be a series of R -submodules of A with each $N_{\lambda+1}/N_\lambda$ being a cyclic R -module and for any limit ordinal λ the R -module N_λ is the union of all R -modules N_μ for $\mu < \lambda$. We use transfinite induction on β to prove that there is a decreasing filtration $\{F_\mu^d\}_{d \in \mathbb{Z}}$ of R -submodules of N_μ for each $\mu \leq \beta$ with the properties that $\bigcup_{d \in \mathbb{Z}} F_\mu^d = N_\mu$ for all $\mu \leq \beta$, if μ is not a limit ordinal $F_\mu^d \cap N_{\mu-1} = F_{\mu-1}^d$, and if μ is a limit ordinal $F_\mu^d = \bigcup_{\lambda < \mu} F_\lambda^d$.

First we consider the case when $\beta = 1$, so $A = N + Ra_0$ for some element a_0 of A .

Let $I = \{\lambda \in R: \lambda a_0 \in N\}$ and $\{U^d\}_{d \in \mathbb{Z}}$ be the decreasing filtration of R -submodules of I given by $U^d = \{\lambda \in I: \lambda a_0 \in F^d\}$. Since $I = \bigcup_d U^d$ and R is Noetherian $I = U^{d_0}$ for some integer d_0 . We define a decreasing filtration $\{U_1^d\}_{d \in \mathbb{Z}}$ of R by $U_1^d = U^d$ for $d \geq d_0$ and $U_1^d = R$ for $d \leq d_0 - 1$. This gives a decreasing filtration $\{F_1^d\}$ of A such that $a \in F_1^d$ if and only if $a = a' + \lambda a_0$ for some a' in $F^d \subseteq N$ and $\lambda \in U_1^d$. If in addition a is an element of N then λ is an element of $I = U^{d_0}$. In this situation if $d_0 \geq d$

then $\lambda a_0 \in U^{d_0} a_0 \subseteq F^{d_0} \subseteq F^d$ and if $d_0 \leq d$ the inclusion $U_1^d a_0 = U^d a_0 \subseteq F^d$ yields as well $\lambda a_0 \in F^d$. Thus $F_1^d \cap N = F^d$ as required.

For the inductive step suppose we have constructed decreasing filtrations $\{F_\mu^d\}_{d \in \mathbb{Z}}$ for all $\mu < \beta$. If β is a limit ordinal define $F_\beta^d = \bigcup_{\mu < \beta} F_\mu^d$ for all $d \in \mathbb{Z}$. In the case when β is not a limit ordinal we use the same as in the case $\beta = 1$ to construct a filtration $\{F_\beta^d\}_{d \in \mathbb{Z}}$ such that $F_\beta^d \cap N_{\beta-1} = F_{\beta-1}^d$ for all $d \in \mathbb{Z}$.

Finally we define F_*^d to be F_β^d for all $d \in \mathbb{Z}$.

3.2. An Element of Åberg's Type

We amend slightly the notation given in Section 3.1. We remind the reader that there are m characters χ_1, \dots, χ_m of Q such that $\chi_i(q) = (q_{\chi_i}, q)$ for all $q \in Q$ and some elements $q_{\chi_1}, \dots, q_{\chi_m}$ in Q , where $Q = \mathbb{Z}^n \subset \mathbb{R}^n$ and (\cdot, \cdot) is the standard inner product in \mathbb{R}^n . Furthermore, q_{χ_m} lies in the subgroup Q' of Q generated by $q_{\chi_1}, q_{\chi_2}, \dots, q_{\chi_{m-1}}$ and $Q = Q' \times Q''$ where Q'' is the orthogonal complement of Q' in Q with respect to the standard inner product in \mathbb{R}^n .

For any character χ_i we define a filtration $\{F_{\chi_i}^d\}_{d \in \mathbb{Z}}$ of $\mathbb{Z}Q_{\chi_i}$ -submodules of $N = \mathbb{Z}Q/P_1 \oplus \dots \oplus \mathbb{Z}Q/P_s$ as follows. Let $A_j = \mathbb{Z}Q/P_j$, $A_{j, \chi_i}^d = (q_{\chi_i}^d \mathbb{Z}Q_{\chi_i} + P_j)/P_j$ and $F_{\chi_i}^d = A_{1, \chi_i}^d \oplus \dots \oplus A_{s, \chi_i}^d$, where $Q_{\chi_i} = \{q \in Q: \chi_i(q) \geq 0\}$. By Lemma 3.2 since the ring $\mathbb{Z}Q_{\chi_i}$ is Noetherian the decreasing filtration $\{F_{\chi_i}^d\}_{d \in \mathbb{Z}}$ of $\mathbb{Z}Q_{\chi_i}$ -submodules of N can be extended to a decreasing filtration of $\mathbb{Z}Q_{\chi_i}$ -submodules of A . We write $\{F_{\chi_i}^d\}_{d \in \mathbb{Z}}$ for the new filtration as well. Finally we define a decreasing filtration $\{F^d\}_{d \in \mathbb{Z}}$ of A by $F^d = \bigcap_{i=1}^m F_{\chi_i}^d$. Since $Q'' \subseteq \text{Ker } \chi_i$ for all $1 \leq i \leq m$, each F^d is a $\mathbb{Z}Q''$ -submodule of A .

We define

$$Q'_k = \{q \in Q': \chi_i(q) \geq -k \chi_i(q_{\chi_i}), 1 \leq i \leq m\}, \quad S_k = Q'_k \setminus Q'_{k-1} \quad (1)$$

for any positive integer k and $Q'_0 = S_0 = \{1_Q\}$. Thus $Q'_k = \bigcup_{j \leq k} S_j$. Denote $(Q'_k)^{-1} = \{q \in Q': q^{-1} \in Q'_k\}$ and $S_k^{-1} = \{q \in Q': q^{-1} \in S_k\}$.

Let $\pi: Q \rightarrow G$ be a lifting of the projection map $G \rightarrow Q$. We define a path γ in $(Q'_k)^{-1}$ from 1_Q to an element q of $(Q'_k)^{-1}$ to be any sequence

$$\left(1_Q, q_{\chi_{i(1)}}^{\epsilon(1)}, q_{\chi_{i(2)}}^{\epsilon(2)} q_{\chi_{i(1)}}^{\epsilon(1)}, \dots, \prod_{r=j}^1 q_{\chi_{i(r)}}^{\epsilon(r)} = q \right)$$

of elements of $(Q'_k)^{-1}$, where j is a positive integer, $\epsilon(r) = \pm 1$, $i(r) \in \{1, 2, \dots, m-1\}$. For any such path γ there is a corresponding sequence

$$\begin{aligned} & (1_G, \pi(q_{\chi_{i(1)}})^{\epsilon(1)}, \pi(q_{\chi_{i(2)}})^{\epsilon(2)} \pi(q_{\chi_{i(1)}})^{\epsilon(1)}, \dots, \\ & \pi(q_{\chi_{i(j)}})^{\epsilon(j)} \pi(q_{\chi_{i(j-1)}})^{\epsilon(j-1)} \dots \pi(q_{\chi_{i(1)}})^{\epsilon(1)}) \end{aligned}$$

of elements of the group G . We write $l(\gamma)$ for the last element of the above sequence and define $W_{k,q}$ to be the set of all elements $l(\gamma)$ when γ runs through all paths in $(Q'_k)^{-1}$ from 1_Q to q . Note that $l(\gamma)$ is an element of G such that $l(\gamma)A = q$ and that

$$W_{k,q} = l(\gamma)W_{k,1_Q}. \quad (2)$$

A similar definition of a path could be given with the modified condition that $i(r) \in \{1, 2, \dots, m\} \setminus \{j\}$ for some $j \leq m$ or even that $i(r) \in \{1, 2, \dots, m\}$. We preferred using in the definition of a path the condition that $i(r) \in \{1, 2, \dots, m-1\}$ because $\{q_{\chi_1}, \dots, q_{\chi_{m-1}}\}$ is a \mathbb{Z} -basis of Q' .

The following lemma is the only place we use the hypothesis that either G is a split extension of A by Q or A is of finite exponent.

LEMMA 3.3. $W_{k,1_Q}$ is a finite subgroup of A .

Proof. Note that any element of $W_{k,1_Q}$ corresponds to a closed path at 1_Q in $(Q'_k)^{-1}$ and the group of closed paths at 1_Q in $(Q'_k)^{-1}$, with group operation the concatenation of paths, is finitely generated. Thus $W_{k,1_Q}$ is a finitely generated subgroup of A . If G is a split extension of A by Q we can assume that π is an embedding of Q in G and hence $W_{k,1_Q}$ contains only the trivial element in A .

If G is non-split then, since by assumption A is a torsion abelian group, $W_{k,1_Q}$ is a finite subgroup of A as required.

We shall prove in Section 3.3 that there exist positive integers k_0 and d such that

$$S_k \subseteq \bigcup_{s \in S_{k+1}} O(s, d) \quad \text{for all } k \geq k_0, \quad (3)$$

where $O(s, d)$ is the set of all elements in Q' contained in the open ball in $\mathbb{R} \otimes Q' \simeq \mathbb{R}^{m-1}$ with radius d and centre s . Furthermore we will show in Section 3.3 that there exist positive integers n_0 and t_0 and some subsets U_{k+t_0} of S_{k+t_0} such that

$$S_k \subseteq \bigcup_{u \in U_{k+t_0}} O(u, n_0) \quad \text{for all } k \geq k_0. \quad (4)$$

Denote by $\alpha_{k,q}$ the sum of the elements of $W_{k,q} \subset G$ in the group algebra $\mathbb{Z}G$. We write the elements of the group algebra $\mathbb{Z}G$ as \mathbb{Z} -linear combinations of $\pi(q_1)\pi(q_2)\cdots\pi(q_j)T^a = T^{(q_1q_2\cdots q_j)^a}\pi(q_1)\pi(q_2)\cdots\pi(q_j)$ for $q_1, q_2, \dots, q_j \in Q$, $a \in A$ (these expressions are not unique). We define

$$\alpha_k = \sum_{q \in (Q'_k)^{-1}} \alpha_{k,q} \prod_{u \in U_{k+t_0}} (T^u - 1) \in \mathbb{Z}G,$$

where $U_{k+t_0} \subseteq Q$ embeds in $N = \mathbb{Z}Q/P_1 \oplus \cdots \oplus \mathbb{Z}Q/P_s \subseteq A$ via the diagonal map. Set

$$\alpha = (\alpha_k)_{k \geq k_1} \in \mathbb{Z}G^{\mathbb{N}}, \quad (5)$$

where k_1 is a sufficiently big positive integer; we will define it at the beginning of Section 3.4.

LEMMA 3.4. *Provided (3) and (4) hold and k_1 is a sufficiently big positive integer the image β of α in $E = (\mathbb{Z}G^{\mathbb{N}})_A$ is Q' -invariant.*

Proof. By (2)

$$\alpha_{k,q} = l(\gamma) \sum_{a \in W_{k,1_Q}} T^a$$

for any path γ in $(Q'_k)^{-1}$ from 1_Q to q . If $q \in (Q'_k)^{-1} \cap q_{\chi_i}^{-1}(Q'_k)^{-1}$ for some i between 1 and $m-1$ then

$$\pi(q_{\chi_i})\alpha_{k,q} = \pi(q_{\chi_i})l(\gamma) \sum_{a \in W_{k,1_Q}} T^a = l(\gamma') \sum_{a \in W_{k,1_Q}} T^a = \alpha_{k,q_{\chi_i}q},$$

where γ' is the concatenation of the path γ from 1_Q to q and the path (q, q_{χ_i}) and both paths are in $(Q'_k)^{-1}$.

Then for every $1 \leq i \leq m-1$ one calculates

$$\begin{aligned} \pi(q_{\chi_i})\alpha_k - \alpha_k &= \sum_{q \in (Q'_k)^{-1}} \pi(q_{\chi_i})\alpha_{k,q} \prod_{u \in U_{k+t_0}} (T^u - 1) \\ &\quad - \sum_{q \in (Q'_k)^{-1}} \alpha_{k,q} \prod_{u \in U_{k+t_0}} (T^u - 1) \\ &= \sum_{q \in (Q'_k)^{-1} \cap q_{\chi_i}^{-1}(Q'_k)^{-1}} \alpha_{k,q_{\chi_i}q} \prod_{u \in U_{k+t_0}} (T^u - 1) \\ &\quad + \sum_{q \in (Q'_k)^{-1} \setminus q_{\chi_i}^{-1}(Q'_k)^{-1}} \pi(q_{\chi_i})\alpha_{k,q} \prod_{u \in U_{k+t_0}} (T^u - 1) \\ &\quad - \sum_{q \in (Q'_k)^{-1}} \alpha_{k,q} \prod_{u \in U_{k+t_0}} (T^u - 1) \\ &= \sum_{q \in (Q'_k)^{-1} \setminus q_{\chi_i}^{-1}(Q'_k)^{-1}} \pi(q_{\chi_i})\alpha_{k,q} \prod_{u \in U_{k+t_0}} (T^u - 1) \\ &\quad - \sum_{q \in (Q'_k)^{-1} \setminus q_{\chi_i}^{-1}(Q'_k)^{-1}} \alpha_{k,q} \prod_{u \in U_{k+t_0}} (T^u - 1). \end{aligned} \quad (6)$$

Consider the element $\pi(q_{\chi_i})\alpha_{k,q}\prod_{u\in U_{k+t_0}}(T^u-1)$ for some $q\in(Q'_k)^{-1}\setminus q_{\chi_i}^{-1}(Q'_k)^{-1}$ and $k\geq k_1$. Then $q^{-1}\in Q'_k\setminus q_{\chi_i}Q'_k\subseteq S_k\cup\ldots\cup S_{k-b}$ for some sufficiently big positive number b not depending on k and by (3) there exists an element $s\in S_k$ such that $q^{-1}\in O(s, bd)$ if $k\geq k_1\geq k_0+b$. By (4) there exists an element u_0 in U_{k+t_0} such that $s\in O(u_0, n_0)$. Then $qu_0\in O(1_Q, n_0+bd)$ and

$$\begin{aligned}\pi(q_{\chi_i})\alpha_{k,q}\prod_{u\in U_{k+t_0}}(T^u-1) &= \prod_{u\in U_{k+t_0}}(T^{q_{\chi_i}q^u}-1)\pi(q_{\chi_i})\alpha_{k,q} \\ &\in \sum_{r\in q_{\chi_i}O(1_Q, n_0+bd)}(T^r-1)\mathbb{Z}G\end{aligned}\quad (7)$$

We remind the reader that the elements of Q are viewed as elements of $N\subseteq A$ via the diagonal map from $\mathbb{Z}Q$ to $N=\mathbb{Z}Q/P_1\oplus\ldots\oplus\mathbb{Z}Q/P_s$.

Now we consider $\alpha_{k,q}\prod_{u\in U_{k+t_0}}(T^u-1)$ for $q\in(Q'_k)^{-1}\setminus q_{\chi_i}(Q'_k)^{-1}$ for some $k\geq k_1$. Then $q^{-1}\in Q'_k\setminus q_{\chi_i}^{-1}Q'_k$ and

$$Q'_k\setminus q_{\chi_i}^{-1}Q'_k\subseteq S_k\cup S_{k-1}\cup\ldots\cup S_{k-b}\quad (8)$$

for some positive integer b not depending on k but probably depending on i . Choosing b sufficiently big we can assume that (8) holds for all $1\leq i\leq m$. Assume further that $k_1\geq b+k_0$ and hence $k\geq b+k_0$. Then $q^{-1}\in S_{k-j}$ for some $0\leq j\leq b$ and by (3) there exists an element s in S_k such that $q^{-1}\in O(s, jd)\subseteq O(s, bd)$. Now by (4) we have an element u_0 in U_{k+t_0} such that $s\in O(u_0, n_0)$ and hence $qu_0\in O(1_Q, bd+n_0)$. Then one calculates

$$\alpha_{k,q}\prod_{u\in U_{k+t_0}}(T^u-1)=\prod_{u\in U_{k+t_0}}(T^{q^u}-1)\alpha_{k,q}\in\sum_{r\in O(1_Q, bd+n_0)}(T^r-1)\mathbb{Z}G\quad (9)$$

By (6), (7), and (9) we deduce that the image β of α in $(\mathbb{Z}G^{\mathbb{N}})_A$ is q_{χ_i} -invariant for $1\leq i\leq m-1$ and hence it is Q' -invariant.

3.3. The Choice of k_0 , t_0 , and U_k

In this section we find sets U_k and positive integers k_0, d, t_0 which satisfy (3) and (4) from Section 3.2.

Let A and B be the subsets of \mathbb{R}^s for some positive integer s . We define $\delta(A, B)$ to be the infimum of all real numbers d such that for any element $a\in A$ there exists an element $b\in B$ with the property that the distance between a and b is not more than d . Note that in general $\delta(A, B)\neq\delta(B, A)$. Later we will need the following simple fact.

LEMMA 3.5. *Let s be a natural number and let l_1, l_2, \dots, l_{s+1} be linear functionals on \mathbb{R}^s such that their convex hull in $\text{Hom}(\mathbb{R}^s, \mathbb{R}) \simeq \mathbb{R}^s$ is an s -simplex with the origin as an internal point. Suppose that for some real numbers $a_i \geq b_i$, $1 \leq i \leq s+1$ the sets $A = \{r \in \mathbb{R}^s \mid l_i(r) \geq a_i \text{ for all } 1 \leq i \leq s+1\}$ and $B = \{r \in \mathbb{R}^s \mid l_i(r) \geq b_i \text{ for all } 1 \leq i \leq s+1\}$ are s -simplexes. Denote the sets of vertices of A and B by A° and B° , respectively, so $A^\circ = \{a_1^\circ, \dots, a_{s+1}^\circ\}$, $B^\circ = \{b_1^\circ, \dots, b_{s+1}^\circ\}$ where $l_i(a_j^\circ) = a_i$ for $1 \leq i \neq j \leq s+1$ and $l_i(b_j^\circ) = b_i$ for all $1 \leq i \neq j \leq s+1$. Then $A \subseteq B$ and*

$$\delta(B, A) \leq \max\{\|a_i^\circ - b_i^\circ\|: 1 \leq i \leq s+1\}. \quad (1)$$

If in addition $0 \in A$ and $B = r_0 A$ for some real number $r_0 > 1$, we have

$$\delta(\overline{B \setminus A}, \delta B) \leq \max\{\|a_i^\circ - b_i^\circ\|: 1 \leq i \leq s+1\}, \quad (2)$$

where δB denotes the boundary of B and $\overline{B \setminus A}$ is the closure of $B \setminus A$.

Remark. Inequality (2) is true without the additional hypothesis but we need it only in the described form.

Proof. (1) Let b be an element of B and $a \neq b$ an element of $A \setminus \delta B$. Then the intersection of the line through a and b and the boundary δB of B contains only two distinct points, say b_1 and b_2 . Without loss of generality we can assume that b is a point between a and b_2 and so $b = ta + (1-t)b_2$ for some $t \in [0, 1)$. Since $b_2 \in \delta B$ there exists some $i \in \{1, 2, \dots, s+1\}$ and non-negative real numbers $\{\mu_j\}_{1 \leq j \neq i \leq s+1}$ such that $b_2 = \sum_{1 \leq j \neq i \leq s+1} \mu_j b_j^\circ$ and $\sum_{1 \leq j \neq i \leq s+1} \mu_j = 1$. We define $c = ta + (1-t)\sum_{1 \leq j \neq i \leq s+1} \mu_j a_j^\circ$. Since A is a convex set c is an element of A and $\|b - c\| = \|(1-t)(b_2 - \sum_{j \neq i} \mu_j a_j^\circ)\| \leq \|\sum_{j \neq i} \mu_j (b_j^\circ - a_j^\circ)\| \leq \max\{\|b_j^\circ - a_j^\circ\|: 1 \leq j \leq s+1\}$.

(2) Let b be an element of $B \setminus A$ and let the ray from 0 through b intersect the boundary δA of A at the point c_1 and the boundary δB of B at the point c_2 . Then there exists some $i \in \{1, 2, \dots, s+1\}$ and non-negative real numbers $\{\mu_j\}_{1 \leq j \neq i \leq s+1}$ such that $\sum_{j \neq i} \mu_j = 1$, $c_1 = \sum_{j \neq i} \mu_j a_j^\circ$, and $c_2 = \sum_{j \neq i} \mu_j b_j^\circ$. Then $\|b - c_2\| \leq \|c_1 - c_2\| = \|\sum_{j \neq i} \mu_j (b_j^\circ - a_j^\circ)\| \leq \max\{\|b_j^\circ - a_j^\circ\|: 1 \leq j \leq s+1\}$.

Then $\delta(\overline{B \setminus A}, \delta B) = \delta(B \setminus A, \delta B) \leq \max\{\|a_i^\circ - b_i^\circ\|: 1 \leq i \leq s+1\}$, as required.

The following lemma is an easy consequence of Lemma 3.5.

LEMMA 3.6. *Let s be a natural number and let l_1, l_2, \dots, l_{s+1} be linear functionals on \mathbb{R}^s such that their convex hull in $\text{Hom}(\mathbb{R}^s, \mathbb{R}) \simeq \mathbb{R}^s$ is an s -simplex containing the origin as an internal point. Let $A_k = \{r \in \mathbb{R}^s \mid l_i(r) \geq a_{i,k}, 1 \leq i \leq s+1\}$ for some real numbers $a_{i,k}$ and $B_k = \{r \in \mathbb{R}^s \mid l_i(r) \geq a_{i,k} - b_i, 1 \leq i \leq s+1\}$ for some positive real numbers b_i , not depending*

on k , be s -simplexes for all $k \in \mathbb{N}$. Then $A_k \subset B_k$ and

$$\text{the set } \{\delta(B_k, A_k)\}_{k \in \mathbb{N}} \text{ is bounded.} \quad (3)$$

If in addition for all k the origin belongs to A_k and $B_k = r_k A_k$ for some real number $r_k > 1$ then

$$\text{the set } \{\delta(\overline{B_k \setminus A_k}, \delta B_k)\}_{k \in \mathbb{N}} \text{ is bounded.} \quad (4)$$

Proof. Let $A_k^o = \{a_{1,k}^o, a_{2,k}^o, \dots, a_{s+1,k}^o\}$ and $B_k^o = \{b_{1,k}^o, b_{2,k}^o, \dots, b_{s+1,k}^o\}$ be the vertices of A_k and B_k , respectively, given by $l_i(a_{j,k}^o) = a_{i,k}$ for $1 \leq i \neq j \leq s+1$ and $l_i(b_{j,k}^o) = a_{i,k} - b_i$ for $1 \leq i \neq j \leq s+1$. Then $l_i(a_{j,k}^o - b_{j,k}^o) = b_i$ for $1 \leq i \neq j \leq s+1$ does not depend on k , so $a_{j,k}^o - b_{j,k}^o$ does not depend on k . By (1) $\delta(B_k, A_k) \leq \max\{\|a_{j,k}^o - b_{j,k}^o\|: 1 \leq j \leq s+1\}$, so (3) follows immediately. Similarly (4) follows from (2).

By the choice of the characters χ_1, \dots, χ_m for any $1 \leq i \leq m$ there exists an integer $j(i)$ such that

$$\text{the ring } B_i = (\mathbb{Z}Q_{\chi_i} + P_{j(i)}) / (q_{\chi_i} \mathbb{Z}Q_{\chi_i} + P_{j(i)}) \text{ is non-trivial.} \quad (5)$$

Let J_i be a maximal ideal in B_i . Then B_i/J_i is a finite field and there exists a positive integer r_i such that

$$(\text{Ker}(\chi_i))^{r_i} = 1 \quad \text{in } B_i/J_i. \quad (6)$$

We define

$$L_{i,k} = \{q \in S_k: \chi_i(q) = -k\chi_i(q_{\chi_i})\} \subset Q',$$

$$W_{i,k} = \{q \in L_{i,k} \mid \chi_j(q) \geq -(k-1)\chi_j(q_{\chi_j}) \text{ for } 1 \leq j \neq i \leq m\},$$

$$U_{i,k} = \{q \in W_{i,k} \mid qq_{\chi_i}^k \in (\text{Ker } \chi_i)^{r_i}\}, \text{ and set } U_k = \bigcup_{i=1}^m U_{i,k}. \quad (7)$$

We define some subsets of $\mathbb{R}^{m-1} = Q' \otimes \mathbb{R}$ by

$$Q_k^* = \{r \in \mathbb{R}^{m-1} \mid (q_{\chi_j}, r) \geq -k\chi_j(q_{\chi_j}) \text{ for all } 1 \leq j \leq m\},$$

$$S_k^* = Q_k^* \setminus Q_{k-1}^*,$$

$$L_{i,k}^* = \{r \in \mathbb{R}^{m-1} \mid (q_{\chi_j}, r) \geq -k\chi_j(q_{\chi_j}) \text{ for all } 1 \leq j \neq i \leq m,$$

$$(q_{\chi_i}, r) = -k\chi_i(q_{\chi_i})\},$$

$$W_{i,k}^* = \{r \in \mathbb{R}^{m-1} \mid (q_{\chi_j}, r) \geq -(k-1)\chi_j(q_{\chi_j}), 1 \leq j \neq i \leq m,$$

$$(q_{\chi_i}, r) = -k\chi_i(q_{\chi_i})\}.$$

Then $W_{i,k} \subset L_{i,k} \subset S_k \subset Q'_k$ are the points with integral coordinates in $W_{i,k}^* \subset L_{i,k}^* \subset S_k^* \subset Q_k'^* \subset \mathbb{R}^{m-1}$, where Q' is identified with \mathbb{Z}^{m-1} in \mathbb{R}^{m-1} .

The following lemma ensures the existence of positive integers k_0 and d satisfying (3) in Section 3.2.

LEMMA 3.7. *There exist positive integers k_0 and d such that*

$$S_k \subseteq \bigcup_{s \in S_{k+1}} O(s, d) \quad \text{for all } k \geq k_0.$$

Proof. By (4) the sets $\{\delta(Q_k^* \setminus Q_{k-1}^*, \delta(Q_k^*))\}_{k \in \mathbb{N}}$ and $\{\delta(\overline{Q_{k+1}^* \setminus Q_k^*}, \delta(Q_{k+1}^*))\}_{k \in \mathbb{N}}$ are bounded. Since the boundary $\delta(Q_k^*)$ of Q_k^* lies in $\overline{Q_{k+1}^*} \setminus Q_k^*$ we can deduce that the set $\{\delta(S_k^*, \delta(Q_{k+1}^*))\}_{k \in \mathbb{N}}$ is bounded as well.

We claim that for sufficiently large k , say $k \geq k_0$, the set $L_{i,k}$ is non-empty and the set $\{\delta(L_{i,k}^*, L_{i,k})\}_{k \geq k_0}$ is bounded. If this is the case then using that $\delta(Q_{k+1}^*) = \bigcup_{i=1}^m L_{i,k+1}^*$ and $\bigcup_{i=1}^m L_{i,k+1} \subseteq S_{k+1}$ we get the set $\{\delta(S_k^*, S_{k+1})\}_{k \geq k_0-1}$ is bounded and S_k is nonempty for $k \geq k_0$. Consequently there exists a positive integer d such that $\delta(S_k, S_{k+1}) < d$ for all $k \geq k_0$, as required in Lemma 3.7.

Define $B_{i,k} = q_{\chi_i}^k L_{i,k}^* \subseteq (\text{Ker } \chi_i \cap Q') \otimes \mathbb{R} \subset Q' \otimes \mathbb{R} = \mathbb{R}^{m-1}$. Since χ_i is a discrete character there exists a positive integer z_i such that any closed $(m-2)$ -ball in $(\text{Ker } \chi_i \cap Q') \otimes \mathbb{R}$ with radius bigger or equal to z_i contains a point from the free abelian group $Q' \cap \text{Ker } \chi_i$. Define $A_{i,k}$ to be the set of elements r in $B_{i,k}$ such that the $(m-2)$ -ball in $(\text{Ker } \chi_i \cap Q') \otimes \mathbb{R}$ with centre r and radius z_i lies in $B_{i,k}$. Since the maximal radius of an $(m-2)$ -ball entirely contained in $L_{i,k}^*$ goes to infinity as k goes to infinity, for sufficiently big k , say $k \geq k_0$, $A_{i,k}$ is an $(m-2)$ -simplex. Note that by (3) the set $\{\delta(B_{i,k}, A_{i,k})\}_{k \geq k_0}$ is bounded. Then for $k \geq k_0$ we have $\delta(L_{i,k}^*, L_{i,k}) = \delta(q_{\chi_i}^k L_{i,k}^*, q_{\chi_i}^k L_{i,k}) \leq \delta(B_{i,k}, A_{i,k}) + \delta(A_{i,k}, q_{\chi_i}^k L_{i,k}) \leq \delta(B_{i,k}, A_{i,k}) + z_i$ and hence $\{\delta(L_{i,k}^*, L_{i,k})\}_{k \geq k_0}$ is bounded, as required.

The following lemma about the filtration $\{F^d\}_{d \in \mathbb{Z}}$ defined at the beginning of Section 3.2 will play a key role in the proof that the image of the element β in $(\mathbb{Z}G^{\mathbb{N}})_{G^n}$ is nontrivial.

LEMMA 3.8. *For sufficiently large positive integer t_0 we have*

$$W_{k,1_Q} \subseteq F^{-k-t_0+1} \quad (8)$$

for all k .

Proof. Let Γ be the set of all closed paths in Q' of the type

$$(1_Q, q_{\chi_i}^{\epsilon(i)}, q_{\chi_i}^{\epsilon(i)} q_{\chi_j}^{\epsilon(j)}, q_{\chi_j}^{\epsilon(j)}, 1_Q),$$

where $1 \leq i \neq j \leq m-1$ and $\epsilon(i), \epsilon(j) \in \{-1, 1\}$. Then $W_{k, 1_Q}$ is an additive subgroup of A contained in the subgroup generated by $l(\gamma_1)^{-1} l(\gamma) l(\gamma_1)$, where γ_1 is a path in $(Q'_k)^{-1}$ starting from 1_Q and $\gamma \in \Gamma$. Since A is a left $\mathbb{Z}Q$ -module via conjugation and all the paths γ_1 lie in $(Q'_k)^{-1}$ we have

$$W_{k, 1_Q} \subseteq \sum_{\gamma \in \Gamma} \mathbb{Z}Q'_k l(\gamma).$$

Via the diagonal map from Q to N the set Q'_k maps to elements in F^{-k} and hence it is sufficient to choose t_0 such that $l(\gamma) \in F^{-t_0+1}$ for all $\gamma \in \Gamma$; this is possible because Γ is a finite set.

The following proposition ensures the existence of natural numbers n_0 and t_0 satisfying (4) from Section 3.2. The statement is geometrically clear but the precise calculation involves several simple geometrical arguments.

PROPOSITION 3.9. *There exist a positive integer t_0 satisfying (8) and a natural number n_0 , depending on t_0 , such that*

$$S_k \subseteq \bigcup_{u \in U_{k+t_0}} O(u, n_0) \quad \text{for all } k \geq k_0.$$

Proof. (1) We claim that there exists a positive integer d_0 such that the set U_k is non-empty for all $k \geq d_0$ and the set $\{\delta(S_k^*, U_k)\}_{k \geq d_0}$ is bounded.

We have seen in the proof of Lemma 3.7 that the set $\{\delta(S_k^*, \bigcup_{i=1}^m L_{i,k}^*)\}_{k \in \mathbb{N}}$ is bounded. Now $W_{i,k}^* \subset L_{i,k}^*$ are $(m-2)$ -simplexes and we can apply (3) to the $(m-2)$ -simplexes $q_{\chi_i}^k W_{i,k}^* \subset q_{\chi_i}^k L_{i,k}^* \subset (\text{Ker } \chi_i \cap Q') \otimes \mathbb{R}$, for $k \in \mathbb{N}$. Since $\delta(L_{i,k}^*, W_{i,k}^*) = \delta(q_{\chi_i}^k L_{i,k}^*, q_{\chi_i}^k W_{i,k}^*)$ we have that the set $\{\delta(L_{i,k}^*, W_{i,k}^*)\}_{k \in \mathbb{N}}$ is bounded.

It remains to prove that for k sufficiently large, say $k \geq d_0$, $U_{i,k}$ is non-empty for all $1 \leq i \leq m$ and the set $\{\delta(W_{i,k}^*, U_{i,k})\}_{1 \leq i \leq m; k \geq d_0}$ is bounded. If this is the case we can deduce that there exists a positive integer n_1 such that

$$U_k \text{ is non-empty and } \delta(S_k^*, U_k) < n_1 \text{ for all } k \geq d_0. \quad (9)$$

We consider $B_{i,k} = q_{\chi_i}^k W_{i,k}^* = \{q_{\chi_i}^k q \mid q \in W_{i,k}^*\}$, an $(m-2)$ -simplex in $(\text{Ker } \chi_i \cap Q') \otimes \mathbb{R}$. Then $q_{\chi_i}^k U_{i,k} = B_{i,k} \cap (\text{Ker } \chi_i \cap Q')^{r_i} \subseteq Q' = \mathbb{Z}^{m-1}$. Since χ_i is a discrete character the subgroup $Q_i = Q' \cap \text{Ker } \chi_i$ of Q' is a

free abelian of rank $m - 2$ and hence there exists a positive integer z_i such that any closed $(m - 2)$ -ball in $Q_i \otimes \mathbb{R} \subset Q' \otimes \mathbb{R}$, with radius bigger than or equal to z_i , contains a point of $(Q_i)^{r_i}$.

Let $A_{i,k}$ be the set of all elements $r \in B_{i,k}$ such that the closed $(m - 2)$ -ball in $Q_i \otimes \mathbb{R}$ with center r and radius z_i lies in $B_{i,k}$; so $A_{i,k}$ is empty, a single point, or an $(m - 2)$ -simplex contained in $B_{i,k}$. Since $W_{i,k}^*$ is an $(m - 2)$ -simplex with the property that the maximal radius of a closed $(m - 2)$ -ball contained in $W_{i,k}^*$ goes to infinity as k goes to infinity, we have that for sufficiently large k , say $k \geq d_0$, $A_{i,k}$ is an $(m - 2)$ -simplex. Then the set $q_{\chi_i}^k U_{i,k}$ is non-empty for all $k \geq d_0$. By (3) we obtain that the set $\{\delta(B_{i,k}, A_{i,k})\}_{k \geq d_0}$ is bounded. Then $\delta(B_{i,k}, q_{\chi_i}^k U_{i,k}) \leq \delta(B_{i,k}, A_{i,k}) + \delta(A_{i,k}, q_{\chi_i}^k U_{i,k}) \leq \delta(B_{i,k}, A_{i,k}) + z_i$ for $k \geq d_0$. Thus

$$\{\delta(W_{i,k}^*, U_{i,k})\}_{k \geq d_0} = \left\{ \delta(q_{\chi_i}^k W_{i,k}^*, q_{\chi_i}^k U_{i,k}) \right\}_{k \geq d_0} \text{ is bounded}$$

as required.

(2) Let t_0 be any positive number satisfying (8) and $t_0 \geq d_0$. Then by Lemma 3.7 we have

$$S_k \subseteq \bigcup_{u \in S_{k+t_0}} O(u, dt_0) \quad \text{for all } k \geq k_0. \quad (10)$$

Finally set $n_0 = n_1 + dt_0$, where n_1 is the positive integer given in (9). Using (9) and (10) we obtain

$$S_k \subseteq \bigcup_{u \in S_{k+t_0}} O(u, dt_0) \subseteq \bigcup_{u \in U_{k+t_0}} O(u, n_0) \quad \text{for all } k \geq k_0$$

as required.

3.4. The Image of β is Non-trivial in $(\mathbb{Z}G^{\mathbb{N}})_{G''}$

Now we summarize. We take k_0 and d to be the integers defined in Lemma 3.7, then define the positive integers n_0 and t_0 and the sets U_{k+t_0} to be the ones given by Proposition 3.9, and finally set $k_1 = b + k_0$, where b is the positive integer introduced in (8), Section 3.2. Let α be the element of $\mathbb{Z}G^{\mathbb{N}}$ defined in (5), Section 3.2 with the chosen positive integers k_1 and t_0 and the sets U_{k+t_0} , $k \in \mathbb{N}$. Then both conditions (3) and (4) from Sections 3.2 hold, so by Lemma 3.4 the image β of α in $(\mathbb{Z}G^{\mathbb{N}})_A$ is Q' -invariant.

We remind the reader that G'' was defined in Section 3.1 (see Proposition 3.1) and finally we prove

THEOREM 3.10. *The image of β in $(\mathbb{Z}G^{\mathbb{N}})_{G''}$ is non-trivial.*

Proof. Assume, to the contrary, that the image of β in $(\mathbb{Z}G^{\mathbb{N}})_{G''}$ is trivial. By Proposition 3.1 there exists an integer d^* such that

$$\alpha_{k,1_Q} \prod_{u \in U_{k+t_0}} (T^u - 1) = 0 \quad \text{in } \mathbb{Z}[A/F^{d^*}].$$

Note that $\alpha_{k,1_Q}$ was defined before Lemma 3.4 and should not be confused with the similar notation used in Proposition 3.1. By (8) in Section 3.3 we have $\alpha_{k,1_Q} = cT^0$ in $\mathbb{Z}[A/F^{-k-t_0+1}]$ where c is the number of the elements in the finite set $W_{k,1_Q}$; we remind the reader that the set $W_{k,1_Q}$ was defined in Section 3.2. For sufficiently large k , say $k \geq k^*$, $F^{d^*} \subseteq F^{-k-t_0+1}$ and so $\alpha_{k,1_Q} \prod_{u \in U_{k+t_0}} (T^u - 1) = 0$ in $\mathbb{Z}[A/F^{-k-t_0+1}]$. Hence

$$\prod_{u \in U_{k+t_0}} (T^u - 1) = 0 \quad \text{in } \mathbb{Z}[A/F^{-k-t_0+1}] \text{ for } k \geq k^*. \quad (1)$$

Let V_{k+t_0} be the abelian subgroup of A/F^{-k-t_0+1} generated by the classes of the elements in U_{k+t_0} ; we remind the reader that the elements of $U_{k+t_0} \subset Q$ embed in $N \subseteq A$ via the diagonal map. Let $V_{i,k+t_0}$ be the subgroup of V_{k+t_0} generated by the classes of the elements in $U_{i,k+t_0}$. We claim that V_{k+t_0} is a direct sum of its subgroups $V_{i,k+t_0}$ for $1 \leq i \leq m$.

Let λ_i be an element of the abelian subgroup of N generated by $U_{i,k+t_0}$, $1 \leq i \leq m$. By (7) in Section 3.3 we have $\chi_j(U_{i,k+t_0}) \geq -(k+t_0-1)\chi_j(q_{\chi_j})$ for $j \neq i$, so $\lambda_i \in F_{\chi_j}^{-k-t_0+1}$ for $1 \leq j \neq i \leq m$. If $\sum_{i=1}^m \lambda_i \in F^{-k-t_0+1} = \cap_{i=1}^m F_{\chi_i}^{-k-t_0+1} \subseteq F_{\chi_j}^{-k-t_0+1}$, since $\lambda_j \in F_{\chi_j}^{-k-t_0+1}$ for all $j \neq i$, we have $\lambda_i \in F_{\chi_j}^{-k-t_0+1}$. Using again $\lambda_i \in F_{\chi_j}^{-k-t_0+1}$ for $1 \leq j \neq i \leq m$ we get $\lambda_i \in \cap_{j=1}^m F_{\chi_j}^{-k-t_0+1} = F^{-k-t_0+1}$; hence V_{k+t_0} is a direct sum of its subgroups $V_{i,k+t_0}$ for $1 \leq i \leq m$.

One can write (1) of the form $\prod_{u \in U_{k+t_0}} (T^u - 1) = 0$ in $\mathbb{Z}[V_{k+t_0}]$; hence there exists i_0 such that

$$\prod_{u \in U_{i_0,k+t_0}} (T^u - 1) = 0 \quad \text{in } \mathbb{Z}[V_{i_0,k+t_0}] \text{ for } k = k^*.$$

Since

$$U_{i_0,k+t_0} \subseteq F_{\chi_j}^{-k-t_0+1} \quad \text{for } j \neq i_0$$

the abelian group $V_{i_0,k+t_0}$ is isomorphic to the abelian subgroup $V'_{i_0,k+t_0}$ of $A/F_{\chi_{i_0}}^{-k-t_0+1}$ generated by the image of $U_{i_0,k+t_0}$ in N under the diagonal map. Thus

$$\prod_{u \in U_{i_0,k+t_0}} (T^u - 1) = 0 \quad \text{in } \mathbb{Z}[V'_{i_0,k+t_0}] \text{ for } k = k^*$$

and

$$V'_{i_0, k+t_0} \text{ embeds in } \bigoplus_{j=1}^s \left(q_{\chi_{i_0}}^{-k-t_0} \mathbb{Z} Q_{\chi_{i_0}} + P_j \right) / \left(q_{\chi_{i_0}}^{-k-t_0+1} \mathbb{Z} Q_{\chi_{i_0}} + P_j \right). \quad (2)$$

We know by (5) in Section 3.3 that the ring $B_{i_0} = (\mathbb{Z} Q_{\chi_{i_0}} + P_{j(i_0)}) / (q_{\chi_{i_0}} \mathbb{Z} Q_{\chi_{i_0}} + P_{j(i_0)})$ is non-trivial. Obviously $B'_{i_0} := (q_{\chi_{i_0}}^{-k-t_0} \mathbb{Z} Q_{\chi_{i_0}} + P_{j(i_0)}) / (q_{\chi_{i_0}}^{-k-t_0+1} \mathbb{Z} Q_{\chi_{i_0}} + P_{j(i_0)})$ is isomorphic to B_{i_0} as a $\mathbb{Z}(\text{Ker } \chi_{i_0})$ -module. By (2) we have

$$\prod_{u \in U_{i_0, k+t_0}} (T^u - 1) = 0 \quad \text{in } \mathbb{Z}[B'_{i_0}] \text{ for } k = k^*$$

and so

$$\prod_{u \in q_{\chi_{i_0}}^{k+t_0} U_{i_0, k+t_0}} (T^u - 1) = 0 \text{ in } \mathbb{Z}[B_{i_0}] \text{ and hence in } \mathbb{Z}[B_{i_0}/J_{i_0}] \text{ for } k = k^*,$$

where J_{i_0} is the ideal of B_{i_0} defined in Section 3.3. By (6) in Section 3.3 all the elements of $(\text{Ker } \chi_{i_0})^{r_{i_0}}$ represent $1 + J_{i_0}$ in B_{i_0}/J_{i_0} and by (7) in Section 3.3 $q_{\chi_{i_0}}^{k+t_0} U_{i_0, k+t_0} \subseteq (\text{Ker } \chi_{i_0})^{r_{i_0}}$, so

$$(T^{1+J_{i_0}} - 1)_{|U_{i_0, k+t_0}|} = 0 \quad \text{in } \mathbb{Z}[B_{i_0}/J_{i_0}] \text{ for } k = k^*.$$

The last statement contradicts the fact that $1 + J_{i_0}$ is a non-trivial element of the finite field B_{i_0}/J_{i_0} and $\mathbb{Z}[B_{i_0}/J_{i_0}]$ does not have nilpotent elements. This completes the proof of Theorem 3.10.

We have just shown that the image of our special element β under the canonical map

$$H^0(Q', (\mathbb{Z}G^{\mathbb{N}})_A) \rightarrow H_0(Q'', (\mathbb{Z}G^{\mathbb{N}})_A)$$

is non-trivial. This implies that G is not of type FP_m because as shown in [1] the above map factors through $H_{m-1}(Q, (\mathbb{Z}G^{\mathbb{N}})_A) \simeq H_{m-1}(G, \mathbb{Z}G^{\mathbb{N}})$ and the latter should be trivial if G is of type FP_m . This finishes the proof of Theorem B.

It is worth noting that the proof of Theorem B can be used to show the result of Bieri and Strebel [5] that FP_2 implies 2-tameness. As before we can assume that A is not 2-tame and then the subgroup Q' of Q defined in Section 3.1 has rank one. The latter is crucial to claim existence of non-trivial Q' -invariant elements in $(\mathbb{Z}G^{\mathbb{N}})_A$ (in the notation of Section 3.2, since the rank of Q' is one, for any non-negative integer k the subgroup $W_{k, 1_Q}$ of A is trivial), which is the main obstacle in the non-split case.

Finally we remark that in the non-split case the condition that \mathcal{A} is of finite exponent was used only in the proof of Lemma 3.3 but the observation that $W_{k,1_Q}$ is finite is vital for the construction of the element α .

ACKNOWLEDGMENTS

We thank Dr. C. J. B. Brookes for introducing us to the topic and the helpful remarks on a preliminary version of the paper and K. U. Bux for sending a preliminary version of his results while this paper was in preparation. This work was supported by a grant from Trinity College, Cambridge.

REFERENCES

1. H. Åberg, Bieri-Strebel valuations (of finite rank), *Proc. London Math. Soc.* (3) **52** (1986), 269–304.
2. R. Bieri, Homological dimension of discrete groups, *Queen Mary College Math. Notes* (1981).
3. R. Bieri and J. Groves, Metabelian groups of type $(FP)_\infty$ are virtually of type FP , *Proc. London Math. Soc.* (3) **45** (1982), 365–384.
4. R. Bieri and J. Groves, The geometry of the set of characters induced by valuations, *J. Reine Angew. Math.* **347** (1984), 168–195.
5. R. Bieri and R. Strebel, Valuations and finitely presented metabelian groups, *Proc. London Math. Soc.* (3) **41** (1980), 439–464.
6. R. Bieri and R. Strebel, A geometric invariant for modules over an abelian group, *J. Reine Angew. Math.* **322** (1981), 170–189.
7. R. Bieri and R. Strebel, “Geometric Invariants for Discrete Groups,” in preparation.
8. N. Bourbaki, “Elements of Mathematics (Commutative Algebra),” Hermann, Paris, 1972.
9. K. S. Brown, Finiteness properties of groups, *J. Pure Appl. Algebra* **44** (1987), 45–75.
10. K. U. Bux, Finiteness properties of certain metabelian arithmetic groups in the function field case, submitted for publication.
11. G. A. Noskov, Bieri–Strebel invariant and homological finiteness properties of metabelian groups, SFB-Preprint 93-028, Universität Bielefeld, 1993.